

# Telemetric monitoring of 24 h intraocular pressure in conscious and freely moving C57BL/6J and CBA/CaJ mice

Ruixia Li, John H.K. Liu

*Hamilton Glaucoma Center and Department of Ophthalmology, University of California, San Diego, La Jolla, CA*

**Purpose:** To study change patterns of 24 h intraocular pressure (IOP) in conscious and freely moving mice using telemetry.

**Methods:** Adult C57BL/6J and CBA/CaJ mice were entrained to a standard 12 h light and 12 h dark cycle. A telemetric pressure transmitter was implanted subcutaneously on the upper back of each light-dark entrained mouse, and the pressure catheter tip was inserted into the vitreous chamber. Broadcasted IOP data were received at 120 Hz. Means of 2 min IOP were recorded every 5 min for 4–13 days to generate the 24 h IOP pattern in each mouse strain. The pattern of IOP in the C57BL/6J strain was also determined under an acute constant dark condition for 24 h.

**Results:** There were distinct patterns of 24 h IOP in the C57BL/6J and CBA/CaJ mouse strains. Under the standard light-dark condition, IOP was higher during the dark period than the light period in both strains. Elevation in IOP from the light period to the dark period was significantly smaller in the CBA/CaJ strain ( $1.6 \pm 1.7$  mmHg, mean  $\pm$  standard deviation (SD),  $n=21$ ) than in the C57BL/6J strain ( $3.4 \pm 2.5$  mmHg,  $n=20$ ). The 24 h IOP pattern in the C57BL/6J strain persisted under an acute constant dark condition ( $n=8$ ).

**Conclusions:** Distinct change patterns of 24 h IOP appeared in these two mouse strains. Although mean IOP during the dark period was significantly higher than the light period in both strains, the magnitudes of light-dark IOP elevation differed. The 24 h IOP change pattern can be driven endogenously in the absence of light.

Daily variation of intraocular pressure (IOP) is important information for the diagnosis and treatment of glaucoma [1, 2]. To determine IOP variation, continuous IOP monitoring under unrestrained conditions offers significant advantages over periodic IOP measurements under restrained conditions. At the present time, no clinical technique is available to monitor human IOP continuously. However, continuous IOP monitoring has been successfully demonstrated in conscious and freely moving laboratory rabbits using telemetry [3-8]. The telemetric pattern of 24 h IOP in rabbits agrees with the 24 h IOP pattern observed in conscious, restrained rabbits using a pneumatonometer [9,10].

Increasing applications of genetic technology for studying IOP [11-13] have raised interest in the daily IOP variation in mice [14-17]. Until recently, change patterns of 24 h IOP in this nocturnally active rodent have been obtained using periodic IOP measurements involving microneedle cannulations of the anterior chamber under general anesthesia [18]. The 24 h IOP change pattern in conscious and freely moving mice is not known. In the present study, telemetry with a miniaturized pressure transmitter was used to study the 24 h IOP patterns in two inbred mouse strains. One was the widely used C57BL/6J strain. The other was the CBA/CaJ strain that may not have had significant IOP variation between

the light and dark periods compared to many other mouse strains including the C57BL/6J strain [14].

## METHODS

Adult C57BL/6J and CBA/CaJ mice (20–25 g) were obtained from the Jackson Laboratory (Bar Harbor, ME) and entrained to a daily 12 h light (6 AM to 6 PM) and 12 h dark cycle for at least two weeks before the experiments. Food and water were freely available, and the housing temperature was constant at 22 °C. Experiments complied with the guidelines of the Institute for Laboratory Animal Research, and the procedures were approved by the Institutional Animal Care and Use Committee.

A battery-powered, telemetric pressure transmitter (Model PA-C20, weight 3.4 g; Data Sciences International, St. Paul, MN) was implanted subcutaneously on the upper back under aseptic conditions. The mouse was anesthetized with intramuscular ketamine (100 mg/kg) and xylazine (10 mg/kg). A midline incision was made to the dorsal neck. Subcutaneous tissues toward the upper back were gently separated to form a packet for placing the pressure transmitter that was immobilized with sutures to the skin. A subcutaneous fistula was created from the dorsal neck to the temporal eyelid for routing the pressure catheter. Guided by a 20 gauge needle bevel, the catheter tip (0.3–0.4 mm external diameter) was inserted halfway between the limbus and the optic nerve bundle into the vitreous chamber. The catheter tip was advanced 2 mm into the vitreous chamber, passing the pupillary midpoint. The eyeball diameter was 3.5–4 mm. Anchoring the distal pressure catheter on the thin sclera was

---

Correspondence to: John H.K. Liu, Ph.D., University of California, San Diego, Department of Ophthalmology, 9500 Gilman Drive, La Jolla, CA, 92093-0946; Phone: (858) 534-7056; FAX: (858) 534-7014; email: [joliu@ucsd.edu](mailto:joliu@ucsd.edu)



persistent ocular inflammation/irritation could make the implanted pressure catheter tip uncomfortable to the host, and frequent eyeball scratching easily broke the setup. These complications led to no verifiable IOP pulsations online that corresponded to the heart beats. The overall success rate for continuous IOP recordings was about 40% (41/98) in the present study. In the successful recordings, there were minor adverse effects of the implantation to the locomotive activities because of an extra load of the pressure transmitter to the bodyweight. However, there was no alteration in the light/sleep and dark/active behavior patterns.

A complete pressure recording in one C57BL/6J mouse is presented in Figure 1. For this mouse, IOP was relatively low and flat shortly after the surgery and there was no identifiable 24 h pattern for approximately 48 h. A consistent 24 h IOP rhythm appeared after three days and up to 13 days, and significant IOP fluctuations occurred among the 2 min IOP means. In most postoperative eyes, the 2 min IOP fluctuations occurred while normal blinking and eye movement were lessened by the implanted pressure catheter. The spontaneous day-to-day fluctuation in IOP such as that on day 4 in Figure 1 could appear anytime during the data collection period.

Figure 2 summarizes the 24 h patterns of hourly pressure averages in the C57BL/6J mouse strain under three experimental conditions. Repeated measures ANOVA




Figure 2. Change pattern of 24 h intraocular pressure (IOP) in C57BL/6J mice. Error bars represent SEM. Repeated measures ANOVA indicated a statistical difference in the hourly IOP averages under the standard light-dark and the acute constant dark conditions but not in the control experiment.

indicated a statistically significant variation in the hourly IOP averages under the standard light-dark and the acute constant dark conditions ( $p < 0.001$ ) but not in the control experiments ( $p = 0.108$ ). Hourly IOP was in general higher during the dark period than during the light period under the standard light-dark condition. There were two time-dependent IOP rises, one at the beginning of the dark period and the other at the transition from the dark period to the light period. The average IOP during the light period was  $15.6 \pm 5.2$  mmHg (mean  $\pm$  SD,  $n = 20$ ), and the average IOP during the dark period was  $19.0 \pm 5.6$  mmHg. The light-dark IOP difference of  $3.4 \pm 2.5$  mmHg was statistically significant ( $p < 0.001$ ). Figure 2 also shows a similar 24 h IOP change pattern under an acute constant dark condition ( $n = 8$ ). In the five control experiments with the pressure catheter tip placed outside the eyeball, the pressure registered at the transmitter was relatively flat throughout the 24-h cycle. The pressure difference of  $0.3 \pm 1.5$  mmHg between the light period ( $2.8 \pm 3.4$  mmHg) and the dark period ( $3.1 \pm 4.1$  mmHg) was not statistically significant.

Twenty-one postoperative CBA/CaJ mice were housed under the standard 12 h light and 12 h dark cycle, and the change pattern of 24 h IOP was determined. A statistically significant variation in the hourly IOP averages occurred ( $p < 0.001$ ; repeated measures ANOVA). The average IOP was  $12.4 \pm 2.0$  mmHg during the light period and  $14.0 \pm 2.5$  mmHg during the dark period. The IOP difference between the light and dark periods ( $1.6 \pm 1.7$  mmHg) was statistically significant ( $p = 0.024$ ). This intra-strain IOP difference between the light and dark periods in the CBA/CaJ mouse was significantly less than the intra-strain IOP difference in the C57BL/6J mouse ( $p = 0.012$ ). Variations in the hourly IOP from the 24 h means in the CBA/CaJ and C57BL/6J mouse strains are presented in Figure 3. In the CBA/CaJ mouse strain, there was only one time-dependent IOP rise at the beginning of the dark period whereas in the C57BL/6J mouse strain, there were two IOP rises during the 24 h period.

## DISCUSSION

Telemetric IOP monitoring in conscious and freely moving mice offers an alternative to the currently available tonometric methods in mice. Telemetric IOP data can be collected continuously from mice in the absence of general anesthesia [18,20] or any other restrained condition [21-25]. The control experiments on C57BL/6J mice showed relatively stable hourly pressure averages in the recording system. The average pressure difference between the light and dark periods was about 0.3 mmHg in these experiments. When the pressure catheter tip was placed in the vitreous chamber, the 24 h IOP pattern was quite different from the control experiment and the light-dark pressure difference was much larger (3.4 mmHg). This rise and fall pattern of 24 h IOP in the C57BL/6J mouse strain persisted for 24 h under an acute constant dark condition, confirming that the 24 h IOP change

pattern was not driven by light perception and was probably driven endogenously by clock genes in the suprachiasmatic nucleus [16]. The 24 h IOP change pattern might subside under a long-term constant dark condition [17], which was not evaluated in the present study due to the two-week limit on the recording time.

The CBA/CaJ mouse strain surprisingly showed a light-dark IOP difference [14]. However, the magnitude of light-dark IOP difference was statistically smaller in the CBA/CaJ strain than in the C57BL/6J strain, which was partially due to the absence of an IOP rise at the end of the dark period. In a previous study on 24 h pattern of total aqueous humor protein concentration, no difference was found between the C57BL/6J and CBA/CaJ mouse strains [26]. These results suggest that the 24 h rhythms of IOP and total protein concentration in the aqueous humor are not closely related parameters in mice.

We found that the light-dark IOP difference was 3.4 mmHg in the C57BL/6J mouse strain, which was close to the observation using the microneedle method under general anesthesia [16]. Thus, the gross light-dark IOP change pattern in conscious and freely moving mice can be approximated using an invasive tonometry under general anesthesia [15-17]. However, IOP telemetry enabled the pressure data to be sampled at a much higher rate than any other tonometric method. Continuous IOP monitoring can readily provide hour-by-hour IOP information as shown in the present study. Consequently, one is able to detect the small difference in the 24 h IOP change patterns between the C57BL/6J and CBA/CaJ mouse strains.




Figure 3. Variations of intraocular pressure (IOP) from the 24 h means in CBA/CaJ and C57BL/6J mice. Error bars represent SEM. Dashed line represents the zero variation from the 24 h IOP mean. A time-dependent IOP rise during the period of 4–7 PM only appeared in the C57BL/6J strain.

The minute-to-minute IOP fluctuation and the day-to-day spontaneous IOP fluctuation in conscious and freely moving mice (Figure 1) were much larger than the IOP fluctuations previously observed by periodic IOP measurements in anesthetized mice [15-17]. As demonstrated in the telemetric IOP monitoring in rabbits [4], restriction of physical activities during a regular tonometry could dampen the real IOP fluctuation. Fluctuation of IOP in a real life situation is probably more than what can be estimated using a non-continuous tonometry in any species. However, whether or not the large IOP fluctuation observed using telemetry in conscious and freely moving mice is free of artifacts cannot be thoroughly evaluated since there are no other publications in the literature for a comparison.

There are limitations of using telemetry to study IOP in mice. Accuracy of each pressure transmitter to the atmosphere pressure was verified as within  $\pm 3$  mmHg of the manufacturer's calibration before and after each experiment. Since the pressure transmitter was designed to monitor a wide range of blood pressure ( $-20$ – $300$  mmHg), a manufacturer's accuracy limit of  $\pm 3$  mmHg was not an issue for monitoring blood pressure. For IOP monitoring, this accuracy in pressure calibration had its consequences despite the reliability of the pressure recording. A distinct change pattern of 24 h IOP for a specific mouse strain can be considered valid, but the absolute IOP values may not be as accurate. This limitation on accuracy affected the IOP level for each mouse studied and the group means of the C57BL/6J and CBA/CaJ strains. The mean pressure in the control experiments was likely affected as well. Therefore, an accurate IOP difference between the CBA/CaJ strain and the C57BL/6J strain may need to be determined using the microneedle method [14]. On the other hand, the rise and fall of 24 h IOP from the 24-h mean in Figure 3 should not be affected significantly by the inaccuracy in the absolute IOP value.

Using the telemetric IOP model for pharmacological experiments in mice has additional limitations. After inserting the pressure catheter tip into the vitreous chamber, the barrier to ocular delivery of a test agent is no longer intact [27]. Bioavailability of a test agent may also be significantly affected by local tissue reactions to the implantation in the mouse eye. Considering these limitations, we conclude that the telemetric IOP monitoring in conscious and freely moving mice is a useful technique with restrictions. The technique is valid if the goal is to study time-dependent IOP change patterns such as the rise and fall during a circadian cycle. The validity is uncertain if the experimental result depends on absolute IOP values or ocular delivery to an intact eyeball.

#### ACKNOWLEDGMENTS

This study was supported by NIH Grant EY07544 to John H.K. Liu.

## REFERENCES

- Liu JH. Circadian rhythm of intraocular pressure. *J Glaucoma* 1998; 7:141-7. [PMID: 9559503]
- Liu JH. Diurnal measurement of intraocular pressure. *J Glaucoma* 2001; 10:S39-41. [PMID: 11890272]
- Schnell CR, Debon C, Percicot CL. Measurement of intraocular pressure by telemetry in conscious, unrestrained rabbits. *Invest Ophthalmol Vis Sci* 1996; 37:958-65. [PMID: 8631639]
- McLaren JW, Brubaker RF, FitzSimon JS. Continuous measurement of intraocular pressure in rabbits by telemetry. *Invest Ophthalmol Vis Sci* 1996; 37:966-75. [PMID: 8631640]
- Percicot CL, Schnell CR, Debon C, Hariton C. Continuous intraocular pressure measurement by telemetry in alpha-chymotrypsin-induced glaucoma model in the rabbit: effects of timolol, dorzolamide, and epinephrine. *J Pharmacol Toxicol Methods* 1996; 36:223-8. [PMID: 9040114]
- Dinslage S, McLaren J, Brubaker R. Intraocular pressure in rabbits by telemetry II: effects of animal handling and drugs. *Invest Ophthalmol Vis Sci* 1998; 39:2485-9. [PMID: 9804159]
- McLaren JW, Bachman LA, Brubaker RF. Comparison of effects of topical ibopamine and epinephrine on the circadian rhythm of intraocular pressure of the rabbit eye as measured by telemetry. *J Ocul Pharmacol Ther* 1999; 15:107-16. [PMID: 10229488]
- Akaishi T, Ishida N, Shimazaki A, Hara H, Kuwayama Y. Continuous monitoring of circadian variations in intraocular pressure by telemetry system throughout a 12-week treatment with timolol maleate in rabbits. *J Ocul Pharmacol Ther* 2005; 21:436-44. [PMID: 16386085]
- Rowland JM, Potter DE, Reiter RJ. Circadian rhythm in intraocular pressure: a rabbit model. *Curr Eye Res* 1981; 1:169-73. [PMID: 7297102]
- Liu JHK, Gallar J, Loving RT. Endogenous circadian rhythm of basal pupil size in rabbits. *Invest Ophthalmol Vis Sci* 1996; 37:2345-9. [PMID: 8843919]
- John SWM, Anderson MG, Smith RS. Mouse genetics: a tool to help unlock the mechanisms of glaucoma. *J Glaucoma* 1999; 8:400-12. [PMID: 10604301]
- Lindsey JD, Weinreb RN. Elevated intraocular pressure and transgenic applications in the mouse. *J Glaucoma* 2005; 14:318-20. [PMID: 15990617]
- Pang I-H, Clark AF. Rodent models for glaucoma retinopathy and optic neuropathy. *J Glaucoma* 2007; 16:483-505. [PMID: 17700292]
- Savinova OV, Sugiyama F, Martin JE, Tomarev SI, Paigen BJ, Smith RS, John SW. Intraocular pressure in genetically distinct mice: an update and strain survey. *BMC Genet* 2001; 2:12. <http://www.biomedcentral.com/1471-2156/2/12> [PMID: 11532192]
- Aihara M, Lindsey JD, Weinreb RN. Twenty-four-hour pattern of mouse intraocular pressure. *Exp Eye Res* 2003; 77:681-6. [PMID: 14609556]
- Maeda A, Tsujiya S, Higashide T, Toida K, Todo T, Ueyama T, Okamura H, Sugiyama K. Circadian intraocular pressure rhythm is generated by clock genes. *Invest Ophthalmol Vis Sci* 2006; 47:4050-2. [PMID: 16936122]
- Sugimoto E, Aihara M, Ota T, Araie M. Effect of light cycle on 24-hour pattern of mouse intraocular pressure. *J Glaucoma* 2006; 15:505-11. [PMID: 17106363]
- John SW, Hagaman JR, MacTaggart TE, Peng L, Smithes O. Intraocular pressure in inbred mouse strains. *Invest Ophthalmol Vis Sci* 1997; 38:249-53. [PMID: 9008647]
- Aihara M, Lindsey JD, Weinreb RN. Episcleral venous pressure of mouse eye and effect of body position. *Curr Eye Res* 2003; 27:355-62. [PMID: 14704919]
- Avila MY, Carré DA, Stone RA, Civan MM. Reliable measurement of mouse intraocular pressure by a servo-null micropipette system. *Invest Ophthalmol Vis Sci* 2001; 42:1841-6. [PMID: 11431452]
- Cohan BE, Bohr DF. Measurement of intraocular pressure in awake mice. *Invest Ophthalmol Vis Sci* 2001; 42:2560-2. [PMID: 11581198]
- Danias J, Kontiola AI, Filippopoulos T, Mittag T. Method for the noninvasive measurement of intraocular pressure in mice. *Invest Ophthalmol Vis Sci* 2003; 44:1138-41. [PMID: 12601041]
- Reitsamer HA, Kiel JW, Harrison JM, Ransom NL, McKinnon SJ. Tonopen measurement of intraocular pressure in mice. *Exp Eye Res* 2004; 78:799-804. [PMID: 15037114]
- Avila MY, Múnera A, Guzman A, Do CW, Wang Z, Stone RA, Civan MM. Noninvasive intraocular pressure measurements in mice by pneumotometry. *Invest Ophthalmol Vis Sci* 2005; 46:3274-80. [PMID: 16123429]
- Wang WH, Millar JC, Pang IH, Wax MB, Clark AF. Noninvasive measurement of rodent intraocular pressure with a rebound tonometer. *Invest Ophthalmol Vis Sci* 2005; 46:4617-21. [PMID: 16303957]
- Zhou LX, Liu JHK. Circadian variation of mouse aqueous humor protein. *Mol Vis* 2006; 12:639-43. <http://www.molvis.org/v12/a71/> [PMID: 16785852]
- Wang Z, Do CW, Avila MY, Stone RA, Jacobson KA, Civan MM. Barrier qualities of the mouse eye to topically applied drugs. *Exp Eye Res* 2007; 85:105-12. [PMID: 17490649]

The print version of this article was created on 18 April 2008. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.