

Suppression of injury-induced epithelial-mesenchymal transition in a mouse lens epithelium lacking tenascin-C

Sai-ichi Tanaka,¹ Takayoshi Sumioka,¹ Norihito Fujita,¹ Ai Kitano,¹ Yuka Okada,¹ Osamu Yamanaka,¹ Kathleen C. Flanders,² Masayasu Miyajima,³ Shizuya Saika¹

¹Department of Ophthalmology, Wakayama Medical University, Wakayama, Japan; ²Laboratory of Cancer Biology of TGF- β Section, Laboratory of Cancer Biology and Genetics, National Cancer Institute, Bethesda, MD; ³Animal Laboratory, Wakayama Medical University, Wakayama, Japan

Purpose: To investigate the role of tenascin-C in epithelial-mesenchymal transition (EMT) of the lens epithelium during wound healing in mice. Tenascin-C is a component of the extracellular matrix in patients having post-operative capsular opacification.

Methods: The crystalline lens was injured by needle puncture in tenascin-C null (KO, n=56) and wild-type (WT, n=56) mice in a C57BL/6 background. The animals were killed at day 2, 5, or 10 post-injury. Immunohistochemistry was employed to detect α -smooth muscle actin (α SMA), a marker of EMT, collagen type I, transforming growth factor β 1 (TGF β 1), phospho-Smad2, phospho-adducin, and phospho-myosin light chain 9 (MLC9). The expression levels of phospho-adducin and phospho-MLC9 were used as markers for the activation of protein kinase C and Rho kinase, respectively.

Results: The expression of tenascin-C was upregulated in WT lens epithelial cells adjacent to the capsular break at day 5. The results showed that injury-induced EMT of the mouse lens epithelium, as evaluated by histology and the expression patterns of α SMA and fibronectin, was attenuated in the absence of tenascin-C. Upregulation of TGF β 1 expression in the epithelium was also inhibited, and loss of tenascin-C attenuated the phosphorylation of Smad2 and adducin in epithelial cells adjacent to the capsular break. The expression of phospho-adducin was suppressed, while the expression level of phospho-MLC9 was unchanged, in the healing epithelium in the absence of tenascin C.

Conclusions: Tenascin-C is required for injury-induced EMT in the mouse lens epithelium. The mechanism behind this might involve impaired activation of cytoplasmic signaling cascades; i.e., TGF β /Smad and protein kinase C-adding signaling, in the absence of tenascin-C.

Upon injury to the anterior capsule, the lens epithelial cells transdifferentiate into fibroblastic cells, which are capable of expressing fibrous extracellular matrix (ECM) components during the healing process. This process is called the epithelial-mesenchymal transition (EMT) [1,2]. However, EMT often leads to the formation of scar tissue (post-operative capsular opacification, PCO), which impairs post-operative vision, rather than the regeneration of normal lens tissue [3-7]. The expression of α -smooth muscle actin (α SMA) and fibrous collagen is characteristic of myofibroblasts [8-10], and these molecules are also established markers of EMT in lens epithelial cells [3,4]. It has been well established that EMT of in vivo lens epithelial cells and other cell types is induced by transforming growth factor β (TGF β) [2,7,11-18]. Although various signaling cascades are involved in gene expression regulation in EMT, the Smad2/3 and Rho kinase cascades, both of which can be activated by TGF β , are the two major pathways.

Various ECM molecules; i.e., laminin, collagen types I and III, lumican, osteopontin, fibronectin, etc., are upregulated in lens epithelial cells post-EMT [8-10]. In addition, the TGF β signaling EMT involved in various cell types is further modulated by ECM molecules expressed by the cells themselves; i.e., fibronectin, lumican, collagen types I and III, fibulin-5, and osteopontin [19-30]. Integrin-derived signaling activated by matrix binding is also required for EMT.

Tenascin-C is a disulfide-bonded hexamer of a matrix component composed of subunits with molecular weights in the range of 120–300 kDa [31,32]. Tenascin-C expression is also upregulated in post-EMT lens epithelial cells in human PCO [33], suggesting a role for tenascin-C in EMT. A similar mechanism has been postulated for the progression/acquisition of invasive characteristics by cancer cells, which is considered to be a type of EMT; tenascin-C is abundantly detected in the invasive margin of cancers, where the cancer cells express an intermediate filament, vimentin, a marker of EMT in neoplastic cells [34,35]. Moreover, it was reported that the absence of tenascin-C suppresses the development of experimental liver fibrosis in mice [36]. The mechanism behind this involved the attenuation of fibroblast-

Correspondence to: Sai-ichi Tanaka, M.D., Department of Ophthalmology, Wakayama Medical University, 811-1 Kimiidera, Wakayama, 641-0012, Japan; Phone: 81-73-447-2300; FAX: 81-73-448-1991; email: s-tanaka@wakayama-med.ac.jp

myofibroblast conversion and presumably EMT of hepatocytes associated with reduced expression of fibrogenic cytokines. This finding prompted us to hypothesize that loss of this molecule also attenuates EMT *in vivo*; i.e., in the lens epithelium post-injury, because both EMT and fibroblast-myofibroblast conversion produce myofibroblasts, the key player in tissue fibrosis formation. To address this question in the present study, we took advantage of the availability of tenascin-C null (KO) mice [37], and showed that injury-induced EMT of the mouse lens epithelium was perturbed in KO mice.

METHODS

Our experiments were approved by the DNA Recombination Experiment Committee and the Animal Care and Use Committee of Wakayama Medical University, Wakayama, Japan and were conducted in accordance with the Association for Research in Vision and Ophthalmology Statement for the Use of Animals in Ophthalmic and Vision Research.

Lens injury in mouse eyes: Lens injury was performed as previously reported [4,17,31]. We employed the protocol for mouse lens capsular injury approved by National Cancer Institute/National Institutes of Health, Bethesda, MD (Laboratory of Cell Regulation and Carcinogenesis). KO mice with a C57Bl/6 background (n=56) and wild-type (+/+, WT, n=56) mice were generally anesthetized with an intraperitoneal (i.p.) injection of pentobarbital sodium (70 mg/kg bodyweight) and were then topically administered oxybuprocaine eye drops [4,17,26]. Our preliminary investigation showed that the incidence of EMT in the lens epithelium post-injury was similar between the WT and heterozygous mice (data not shown). A small corneal incision was made in the central anterior capsule of one eye (right) with a 26 G hypodermic needle through a corneal incision after topical application of mydriatics, as previously reported [4, 17,26]. In brief, the central anterior lens capsule was pierced once with the blade of the 26 gauge needle. The depth of puncture from the corneal surface was approximately 300 μm , which was about one quarter of the length of the needle blade. The animals that accidentally received a deeper lens injury were not included in the experiment. After instillation of ofloxacin ointment, the mice were allowed to heal for 2, 5, or 10 days. The number of eye samples were 20/20, 21/21, and 21/21 for the WT/KO mice at 2, 5, and 10 days, respectively. Most of the enucleated globes (16/16, 17/17, and 17/17 for the WT/KO mice at 2, 5, and 10 days, respectively) were fixed and embedded in paraffin, and the rest were embedded in optimal cutting temperature (OCT) compound for cryosectioning, as previously reported [4,17, 26].

Histology and immunohistochemistry: Deparaffinized sections and fixed cultured cells were processed for immunohistochemistry as previously reported [4,17,26]. The antibodies used were anti-tenascin C antibody (1:100 in PBS;

Chemicon, Temecura, CA), mouse monoclonal anti- α SMA antibody (1:100 in PBS; Neomarker, Fremont, CA), goat polyclonal anti-fibronectin antibody (1:100 in PBS; Southern Biotechnology, Birmingham, AL), goat-polyclonal anti-phospho-Smad2 antibody (1:100 in PBS; Chemicon), goat polyclonal anti-phospho-adducin (Ser 726) antibody (1:200 in PBS; Santa Cruz Biotechnology, Santa Cruz, CA), goat polyclonal anti-TGF β 1 antibody (1:200 in PBS; Santa Cruz Biotechnology), goat polyclonal anti-TGF β 2 antibody (1:200 in PBS; Santa Cruz Biotechnology), goat polyclonal anti-TGF β receptor type I (TGF β -RI) antibody (1: 200 in PBS, Santa Cruz Biotechnology), and goat polyclonal anti-TGF β receptor type II (TGF β -RII) antibody (1:200 in PBS; Santa Cruz Biotechnology). After being allowed to react with the secondary peroxidase-conjugated antibody and washed in PBS, the antibody complex was visualized using 3,3'-diaminobenzidine. Immunostaining for TGF β 1 was performed as previously reported [17]. Activation of the protein kinase C cascade was assayed by immuno-detection of phospho-adducin (Ser 726). Activation of Rho kinase signaling was evaluated by immuno-detection of phospho (Thr 18/Ser 19)-myosin light chain 9 in cryosections (7 μm in thickness) fixed in cold acetone. After nuclear counterstaining with methyl green, the specimens were observed under regular light microscopy.

RESULTS

Immunohistochemistry for tenascin-C: As tenascin-C expression is reportedly upregulated in human lens epithelial cells during PCO, we first examined whether mouse lens epithelial cells express tenascin-C post-injury using immunohistochemistry. No immunoreactivity for tenascin-C was detected in the uninjured lens epithelia or the injured lens epithelia at day 2 post-injury, but tenascin-C immunoreactivity was observed in the lens epithelium cells adjacent to the break in the anterior capsule at day 5 post-injury (Figure 1). The KO lens epithelium did not express tenascin-C post-injury (data not shown).

Histology of injured mouse lenses: We examined the effects of endogenous tenascin-C on injury-induced EMT of the lens epithelium using HE staining. The histology of the HE-stained tissues showed a marked alteration in the morphology of the cells at the injury site in the WT mice (Figure 2). The lens epithelial cells of the uninjured lens and those at the capsular break at day 2 post-injury maintained a non-elongated morphology in both the WT (Figure 2A and insert) and KO mice (Figure 2B and insert). At day 5 (Figure 2C and insert) and 10 (Figure 2E and insert), the WT cells exhibited an elongated, fibroblast-like morphology in all the specimens examined, suggesting that EMT had occurred. At these time points, the lens capsule adjacent to the break was folded, presumably by contraction of the myofibroblasts in the WT tissues, while it did not appear to be folded or contracted in the KO tissues. On the other hand, the KO lens epithelial cells




Figure 1. Expression of tenascin-C protein in the injured mouse lens epithelium. Neither the epithelium of an uninjured lens (A) nor that of an injured lens at day 2 post-injury (B, asterisk) was labeled for tenascin-C. C: Immunoreactivity for tenascin C was observed in the lens epithelium (asterisk) adjacent to the break in the anterior capsule at day 5 post-injury. The arrows indicate the edge of the break in the anterior capsule. Bar, 10 μ m.




Figure 2. Histology of the edge of the break in the anterior lens capsule in wild-type (WT) and tenascin-C null (KO) mice. At day 2, the lens epithelial cells around the capsular break (arrows) exhibited an epithelial-like morphology in both the WT (A and insert; asterisk) and KO (B and insert; asterisk) mice. At day 5, the epithelial cells had formed a multilayer at the edge of the broken capsule in both the WT (C) and KO (D) mice. The wound indicated by the arrows was filled with cells in the WT mice (C), while that in the KO mouse lens remained open (D). At a higher magnification, the WT cells (C, insert) showed an elongated fibroblast-like shape (asterisk), while those of the KO lens (D, insert) exhibited an epithelial-like morphology (asterisk). At day 10, a multi-cellular layer had formed beneath the anterior capsule in both the WT (E) and KO (F) mice. At a higher magnification, the cells in both groups (E and F; inserts) exhibited an elongated fibroblast-like shape. Inserts of A to F show the higher magnification images of the boxed areas in A through F. AC, anterior lens capsule, Bar, 10 μ m (A through F), or 25 μ m for each of the inserts.

were of epithelial shape, even at day 5, in all specimens examined (Figure 2D and insert). Finally, the KO cells exhibited a fibroblastic, elongated morphology at day 10 (Figure 2F and insert).

Immunohistochemistry for EMT-related components: HE histology suggested that loss of tenascin-C attenuated the EMT process in the lens epithelium. Therefore, we conducted immunodetection of α SMA, fibronectin, and TGF β 1 to confirm this hypothesis. Our previous papers demonstrated the upregulation of the lens epithelium α SMA protein level at

day 5 post-injury. Both the WT and KO epithelial cells of the uninjured lenses and those of the injured lenses from each mouse type were negative for α SMA at day 2 (not shown). The multi-layered fibroblast-like lens cells observed at the capsular break in the WT mice were markedly labeled for α SMA at day 5 (Figure 3A), whereas the epithelial-shaped lens cells in 15 of 17 KO mice were negative for α SMA at this time point (Figure 3B). At day 10, the majority of the cells beneath the broken capsule were labeled with anti- α SMA antibody both in the WT and the KO lenses (Figure 3C,D). As




Figure 3. Expression pattern of α -smooth muscle actin (α SMA) in lens epithelial cells of injured lenses. The multicellular layer of epithelial cells at the edge of the broken capsule was strongly labeled with anti- α SMA antibody in a wild-type mouse at days 5 (A, asterisk) and 10 (C, asterisk). On the other hand, in a tenascin-C null (KO) mouse almost none of the epithelial cells in the multicellular layer were stained for α SMA at day 5 (B). At day 10, the majority of the cells were labeled in the KO mouse (D, asterisk). Arrows, edges of the injured anterior capsule. AC, anterior capsule; Bar, 10 μ m.

shown by HE histology, at day 5 an area of the lens capsule adjacent to the break was folded, presumably by the contraction of myofibroblasts in the WT tissues, while folding or contraction was seen in the KO tissues.

Fibronectin is a major adhesive glycoprotein involved in wound healing. Immunostaining clearly showed that fibronectin protein expression in the lens epithelial cells of the injured lens was inhibited in the KO mice. At day 2, no fibronectin was detected in the injured lens epithelia of the WT or KO mice (not shown). At day 5, loss of tenascin-C suppressed the upregulation of fibronectin expression in the injured lens epithelium; lens cells of a fibroblastic morphology (that were labeled for α SMA) were labeled for fibronectin in WT mice (Figure 4A), although the cells in the KO mice were negative (Figure 4B). At day 10, the lens epithelia in the injured lenses of both the WT and KO mice were positive for fibronectin (Figure 4C,D).

Although we previously reported that the most important type of TGF β for the activation of Smad signaling in the injured mouse lens epithelium is TGF β 2 derived from the aqueous humor [4], there is a possibility that autocrine TGF β 1 also affects these cells. The present immunohistochemical analysis did not detect TGF β 1 in the uninjured lens epithelium, as previously reported (data not shown) [17]. TGF β 1 expression (Figure 5A) was upregulated

in the EMT lens epithelium in the WT mice, while this upregulation was quite minimal in the KO mice at day 2. At day 5, the anterior lens epithelium adjacent to the capsular break was labeled for TGF β 1 in both the WT and KO mice (Figure 5C,D). TGF β 2 was detected in the whole the lens epithelium of the uninjured and healing lenses in both the WT and KO mice (data not shown). TGF β 2 was constitutively expressed in the lens epithelium, and its expression pattern in the healing epithelium was not affected by the loss of TNC (data not shown).

Regarding the TGF β receptors, we examined the expression patterns of TGF β -RI and TGF β -RII in the injured lens epithelium using immunohistochemistry. Both the WT and KO lens uninjured lens epithelia were very faintly labeled with antibodies against TGF β -RI and TGF β -RII (data not shown). At day 5 post-injury, multilayered lens epithelial cells in the injured lenses were markedly labeled with anti-TGF β -RI in the WT mice, while its immunoreactivity was quite faint in the cells from the KO mice (Figure 6A,B). At day 10, the cells in the multilayer beneath the anterior lens capsule were labeled for TGF β -RI in both the WT and KO mice (Figure 6C,D). At day 5 post-injury, the multilayered lens epithelial cells in the injured lenses were markedly labeled with anti-TGF β -RII in the WT mice, while its immunoreactivity was quite faint in the cells from the KO mice (Figure 6E,F). Even at day 10, the immunoreactivity for TGF β -RII was more




Figure 4. Expression pattern of fibronectin in injured lenses. At days 5 (A) and 10 (C), immunoreactivity for fibronectin was detected in the multicellular layer formed at the site of capsular injury in a wild-type mouse (B), whereas no such immunoreactivity was observed in the multicellular layer at day 5 in an injured tenascin-C null lens, although it was readily detected at day 10 (D). Arrows, edge of the injured anterior capsule. AC, anterior capsule; Bar, 10 μ m.

prominent in the WT cells than in the KO cells (Figure 6G,H). TGF β -related signaling mediators. As for the TGF β -related signaling cascades, we examined whether Smad2 and adducin were phosphorylated in the epithelia examined. Adducin is a membrane cytoskeletal protein that is phosphorylated in association with the activation of multiple signaling pathways such as the Rho kinase and protein kinase C pathways [38].

The uninjured lens epithelia of the WT and KO mice were negative for phospho-Smad2 (Figure 7A,B). At day 2, lens epithelial cells containing nuclei labeled for phospho-Smad2 were observed adjacent to the break in the anterior capsule in the WT mice (Figure 7C), whereas the epithelial cells in the KO mice were not labeled (Figure 7D). At day 5, many cells with phospho-Smad2-positive nuclei were observed around the capsular break in the WT mice (Figure 7E). At this time point, the majority of the lens epithelial cells in the cell multi-layer that had formed adjacent to the break in the KO mice were negative for phospho-Smad2 (Figure 7F). At day 10, the cells around the break in the anterior capsule were not labeled for phospho-Smad2 in either the WT or KO mice (Figure 7G,H).

We next immunostained phospho-adducin.38 The uninjured lens epithelia of the WT and KO mice were negative for phospho-adducin (data not shown). At day 2, lens epithelial cells labeled for phospho-adducin were seen in the cytoplasm adjacent to the break in the anterior capsule in the WT mice (Figure 8A), whereas no epithelial cells were labeled for phospho-adducin in the KO mice (Figure 8B). At day 5,

the majority of the lens epithelial cells in the cell multi-layer that had formed adjacent to the break in the KO mice were still negative for phospho-adducin (Figure 8D). At this time point, many elongated cells containing phospho-adducin-positive cytoplasm were observed around the capsular break in the WT mice (Figure 8C). At day 10, the cells around the break in the anterior capsule were not labeled for phospho-adducin in either the WT or KO mice (Figure 8G,H).

We then evaluated the activation of Rho kinase signaling by immuno-detection of phospho (Thr 18/Ser 19)-myosin light chain 9. Phosphorylated-myosin light chain 9 was faintly detected as early as day 2 in the lens epithelial cells of the injured lenses of both the WT and KO mice and was markedly observed thereafter in these cells in both genotypes (Figure 9). The levels of immunoreactivity in the lens epithelial cells were similar in both genotypes.

DISCUSSION

The present study was undertaken to elucidate whether tenascin-C plays a significant role in injury-induced EMT of the lens epithelium using tenascin-C-deficient mice. Here, we show that the absence of tenascin-C perturbs injury-induced EMT in mice *in vivo*. The present study clearly answered this question because the myofibroblasts in healing lens must be derived from the cells inside the lens, e. g., the lens epithelium, without contamination of fibroblastic cells from outside the lens. Since HE histology strongly suggested the impairment of injury-induced EMT in the lens epithelium, we conducted




Figure 5. Expression pattern of transforming growth factor β 1 (TGF β 1) in injured lenses. At days 2 (A) and 5 (C), immunoreactivity for TGF β 1 was detected in the lens epithelial cells adjacent to the site of capsular injury in a wild-type mouse. Although no such immunoreactivity was observed in the multicellular layer at day 2 in an injured tenascin-C null lens, it was readily detected at day 10 (D). Arrows, edge of the injured anterior capsule. AC, anterior capsule; Bar, 10 μ m.




Figure 6. Expression patterns of transforming growth factor β -receptor type I and type II in an injured lens epithelium. At day 5 post-injury, multilayered lens epithelial cells in an injured lens were markedly labeled with anti-TGF β -RI in a wild type (WT; A) mouse, while its immunoreactivity was quite faint in the cells of a tenascin C null (KO; B) mouse. At day 10, the cells of the multilayer beneath the anterior lens capsule were labeled for TGF β -RI in both the WT and KO mice (C, D). At day 5 post-injury, the multilayered lens epithelial cells in an injured lens were markedly labeled with anti-TGF β -RII in a WT mouse (E), while its immunoreactivity was quite faint in the cells of a KO mouse (F). Even at day 10, immunoreactivity for TGF β -RII was more prominent in the WT cells than in the KO cells (G, H); Bar, 10 μ m.

immunohistochemistry and detected reduced expression of α SMA (relative to that in the WT lens), a myofibroblast marker, in lens cells in the KO mice at day 5 post-injury. A similar phenomenon was reported previously; i.e., that tenascin-C is specifically expressed at the invasive fronts of cancers, where neoplastic cells acquire a migratory characteristic during EMT [34,35].

The exact mechanism behind this phenomenon remains to be investigated. We previously reported that TGF β /Smad signaling is essential for the occurrence of EMT in injured mouse lenses [4,17,26]. The present immunohistochemical

analysis showed that the activation of the Smad signal was severely suppressed in the epithelial cells of injured KO mouse lenses compared with that in the epithelial cells of injured WT mouse lenses. TGF β 2, but not TGF β 1, predominates in the aqueous humor [39-41]. We also previously reported that TGF β 1 expression is upregulated in the injured lens epithelium in association with the progression of EMT. On the other hand, blocking TGF β 2, but not TGF β 1, in the aqueous humor abrogates Smad signaling in the injured mouse lens epithelium [17], suggesting that the upregulation of TGF β 1 expression in the epithelia upon injury




Figure 7. Expression patterns of phospho-Smad2 in the lens epithelia of injured lenses. Uninjured lens epithelial cells beneath the anterior capsule were not labeled for phospho-Smad2 in either the wild-type (WT; **A**) or tenascin-C null (KO; **B**) mice. At day 2, the nuclei of some epithelial cells were labeled for phospho-Smad2 in an injured WT lens (**C**, arrows). On the other hand, in a KO injured lens the nuclei of the epithelial cells at the leading edge of the cellular multi-layer were not labeled for phospho-Smad2 (**D**). At day 5, many epithelial cells were founded to contain nuclei that were positive for phospho-Smad2 in the multicellular layer at the capsular break site in a WT mouse (**E**, arrows), whereas no phospho-Smad2 was detected in the majority of epithelial cells in multicellular layer in an injured KO lens (**F**). At day 10, the cells around the break in the anterior capsule were not labeled for phospho-Smad2 in either the WT (**G**) or KO (**H**) mice. Dotted lines, border between the multilayered cells and the lens cortex, AC, anterior capsule; Bar, 20 μ m.

might not play a significant role in the activation of Smad signaling. Thus, in the present study, we examined the expression of TGF β 1 in the injured lens epithelium as a marker of the progression of EMT and showed that the upregulation of TGF β 1 expression in the epithelial cells of the injured lens was also attenuated in the absence of tenascin-C. The expression of TGF β 2, as detected by immunohistochemistry, in the lens epithelium was not affected by the absence TNC (data not shown).

Multiple signaling cascades; i.e., the protein kinase C and Rho kinase cascades, are reportedly required for the EMT and

the expression of fibrogenic components, but TGF β and Smad signaling are considered to be essential. Adducin is a membrane-associated actin-binding protein that is phosphorylated by protein kinase C at its COOH-terminal region. Protein kinase C is also involved in cell migration by modulating the morphology of the F-actin cytoskeleton [38, 42]. We therefore examined whether the pattern of the COOH-terminal (Ser 726) phosphorylation of adducin was altered by the loss of tenascin C in the injured lens epithelium. Our results showed that the phosphorylation of adducin in the injured lens epithelium was greatly reduced in the KO mice




Figure 8. Expression patterns of phospho-adducin in the lens epithelia of injured lenses. At day 2, many lens epithelial cells were labeled for phospho-adducin in an injured wild-type (WT) lens (A, asterisk). On the other hand, in a tenascin-C null (KO) injured lens the epithelial cells at the leading edge were not labeled for phospho-adducin (B). At day 5, many epithelial cells were positive for phospho-adducin in the multicellular layer at the capsular break site in a WT mouse (C, asterisk), although its immunoreactivity was weaker than that in the cells at day 2, whereas phospho-adducin was only faintly detected in the majority of epithelial cells (star) in an injured KO lens at day 5 (D). At day 10, the cells around the break in the anterior capsule were not labeled for phospho-adducin in either WT (E) or KO (F) mice; Bar, 10 μ m.

compared with the WT mice, suggesting that the downregulation of protein kinase C activity in the absence of tenascin C contributes to the attenuation of lens cell EMT in addition to the attenuation of Smad signaling. Rho kinase is another signal that is reportedly essential for EMT in many cell types including the lens epithelium [43,44]. We therefore evaluated the activation of Rho kinase signaling via immunodetection of phospho (Thr 18/Ser 19)-myosin light chain 9. Phosphorylated-myosin light chain 9 was faintly detected as early as day 2 in the lens epithelial cells of the injured lenses

of both the WT and KO mice and was markedly observed thereafter in these cells, suggesting that the absence of tenascin C does not affect Rho kinase activity.

We also examined the expression of TGF β receptors in the injured lens epithelium to understand the susceptibility of the cells to ligands. Immunohistochemistry showed that lens epithelial cells upregulated the expression of TGF β -RI and RII as early as day 5 during the healing process and the progression of EMT in the WT mice. On the other hand, as for the expression pattern of α SMA, the upregulation of




Figure 9. Expression patterns of phospho (Thr 18/Ser 19)-myosin light chain 9 in lens epithelia of injured lenses. At day 2, many lens epithelial cells were faintly labeled for phospho-myosin light chain 9 in both an injured wild-type (WT) and a tenascin-C null (KO) injured lens epithelium (A, B, arrows). At day 5, both the WT and KO cells were obviously immunolabeled for phospho-myosin light chain 9 (C, D, arrows). AC, anterior epithelium; Bar, 10 μ m.

TGF β receptor expression was delayed, negative at day 5, and then positive at day 10 in the KO mice. Nevertheless, it is still unclear whether the delayed EMT seen in the KO mice can be ascribed to the suppression of the upregulation of TGF β receptor expression in the absence of tenascin-C. There is a possibility that delayed upregulation of TGF β receptor expression in KO mice is responsible for the impairment of the phosphorylation of Smad2 and adducin in KO cells. On the other hand, another possible mechanism of the impairment of these signaling transmitters is that the impairment of integrin-derived signals caused by the loss of tenascin-C affects TGF β -derived signaling via signaling crosstalk.

It was reported that in the stroma of the eye cornea, injury-induced expression of fibronectin, matrix component essential for tissue repair, is suppressed in the absence of tenascin-C [45]. In addition, integrin-mediated signaling is reportedly required for fibroblast-myofibroblast conversion and EMT in tissue fibrosis. In the present study, reduced expression of fibronectin was also observed in the injured lens epithelia of KO mice, suggesting that impairment of fibronectin-integrin signaling is also involved in the suppression of lens cell EMT in KO mice.

Exploring the mechanism of EMT in the lens epithelium post-injury is necessary to aid our understanding of the pathogenesis of capsular opacification post-cataract surgery. We reported that anti-Smad molecules inhibit EMT in the lens epithelium and suggested their clinical efficacy in the prevention of PCO [46,47]. However, the present study together with previous reports show the important roles of the ECM in the pathogenesis of PCO. Further study is needed to elucidate the matrix component signaling systems that are required for the occurrence of lens cell EMT.

ACKNOWLEDGMENTS

This study was supported by Grants from the Ministry of Education, Science, Sports, and Culture of Japan (C19592036); Mitsui Foundation; Mochida Memorial Foundation; Takeda Science Foundation; and Uehara Foundation.

REFERENCES

- Hay ED, Zuk A. Transformations between epithelium and mesenchyme: normal, pathological, and experimentally induced. *Am J Kidney Dis* 1995; 26:678-90. [PMID: 7573028]
- Roberts AB, Tian F, Byfield SD, Stuelten C, Ooshima A, Saika S, Flanders KC. Smad3 is key to TGF- β -mediated epithelial-to-mesenchymal transition, fibrosis, tumor suppression and metastasis. *Cytokine Growth Factor Rev* 2006; 17:19-27. [PMID: 16290023]
- Saika S, Kawashima Y, Miyamoto T, Tanaka SI, Ohmi S, Minamide A, Yamanaka O, Ohnishi Y, Ooshima A, Yamanaka A. Immunolocalization of prolyl 4-hydroxylase subunits, α -smooth muscle actin, and extracellular matrix components in human lens capsules with lens implants. *Exp Eye Res* 1998; 66:283-94. [PMID: 9533856]
- Saika S, Okada Y, Miyamoto T, Ohnishi Y, Ooshima A, McAvoy JW. Smad translocation and growth suppression in lens epithelial cells by endogenous TGF β 2 during wound repair. *Exp Eye Res* 2001; 72:679-86. [PMID: 11384156]
- Meacock WR, Spalton DJ, Stanford MR. Role of cytokines in the pathogenesis of posterior capsule opacification. *Br J Ophthalmol* 2000; 84:332-6. [PMID: 10684849]
- Wormstone IM. Posterior capsular opacification: a cell biological perspective. *Exp Eye Res* 2002; 74:337-47. [PMID: 12014915]
- Saika S. Relationship between posterior capsule opacification and intraocular lens biocompatibility. *Prog Retin Eye Res* 2004; 23:283-305. [PMID: 15177204]
- Serini G, Bochaton-Piallat M-L, Ropraz P, Geinoz A, Borsi A, Zardi L, Gabbiani G. The fibronectin domain ED-A is crucial for myofibroblastic phenotype induction by transforming growth factor- β 1. *J Cell Biol* 1998; 142:873-81. [PMID: 9700173]
- Serini G, Gabbiani G. Mechanism of myofibroblast activity and phenotypic modulation. *Exp Cell Res* 1999; 250:273-83. [PMID: 10413583]
- Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA. Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol* 2002; 3:349-63. [PMID: 11988769]
- Saika S, Miyamoto T, Tanaka T, Ishida I, Ohnishi Y, Ooshima A. Latent TGF β binding protein-1 and fibrillin-1 in human

- capsular opacification and in cultured lens epithelial cells. *Br J Ophthalmol* 2001; 85:1362-6. [PMID: 11673307]
12. Miettinen PJ, Ebner R, Lopez AR, Derynck R. TGF- β induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors. *J Cell Biol* 1994; 127:2021-36. [PMID: 7806579]
 13. Srinivasan Y, Lovicu FJ, Overbeek PA. Lens specific expression of transforming growth factor β 1 in transgenic mice causes anterior subcapsular cataracts. *J Clin Invest* 1998; 101:625-34. [PMID: 9449696]
 14. Lee EH, Joo C-K. Role of transforming growth factor- β in transdifferentiation and fibrosis of lens epithelial cells. *Invest Ophthalmol Vis Sci* 1999; 40:2025-32. [PMID: 10440257]
 15. Oft M, Akhurst RJ, Balmain A. Metastasis is driven by sequential elevation of H-ras and Smad2 levels. *Nat Cell Biol* 2002; 4:487-94. [PMID: 12105419]
 16. Wormstone IM, Tamiya S, Anderson I, Duncan G. TGF- β 2-induced matrix modification and cell transdifferentiation in the human lens capsular bag. *Invest Ophthalmol Vis Sci* 2002; 43:2301-8. [PMID: 12091431]
 17. Saika S, Kono-Saika S, Ohnishi Y, Sato M, Muragaki Y, Ooshima A, Flanders KC, Yoo J, Anzano M, Liu CY, Kao WW, Roberts AB. Smad3 signaling is required for epithelial-mesenchymal transition of lens epithelium after injury. *Am J Pathol* 2004; 164:651-63. [PMID: 14742269]
 18. Saika S. TGF β pathobiology in the eye. *Lab Invest* 2006; 86:106-15. [PMID: 16341020]
 19. Hay ED. Extracellular matrix alters epithelial differentiation. *Curr Opin Cell Biol* 1989; 14:455-7.
 20. Zuk A, Hay ED. Expression of β 1 integrins changes during transformation of avian lens epithelium to mesenchyme in collagen gels. *Dev Dyn* 1994; 201:378-93. [PMID: 7534501]
 21. Saika S, Miyamoto T, Tanaka S, Tanaka T, Ishida I, Ohnishi Y, Ooshima A, Ishiwata T, Asano G, Chikama T, Shirai A, Liu CY, Kao CW, Kao WW. Response of lens epithelial cells to injury: role of lumican in epithelial-mesenchymal transition. *Invest Ophthalmol Vis Sci* 2003; 44:2094-102. [PMID: 12714648]
 22. Zeisberg M, Bonner G, Maeshima Y, Colorad P, Muller GA, Strutz F, Kalluri R. Collagen composition and assembly regulates epithelial-mesenchymal transdifferentiation. *Am J Pathol* 2001; 159:1313-21. [PMID: 11583959]
 23. Denhardt DT, Noda M, O'Regan AW, Pavlin D, Berman JS. Osteopontin as a means to cope with environmental insults: regulation of inflammation, tissue remodeling, and cell survival. *J Clin Invest* 2001; 107:1055-61. [PMID: 11342566]
 24. Rangaswami H, Bulbule A, Kundu GC. Osteopontin: role in cell signaling and cancer progression. *Trends Cell Biol* 2006; 16:79-87. [PMID: 16406521]
 25. Saika S, Miyamoto T, Ishida I, Ohnishi Y, Ooshima A. Osteopontin: a component of matrix in capsular opacification and subcapsular cataract. *Invest Ophthalmol Vis Sci* 2003; 44:1622-8. [PMID: 12657601]
 26. Saika S, Shirai K, Yamanaka O, Miyazaki K, Okada Y, Kitano A, Flanders KC, Kon S, Uede T, Kao WW, Rittling SR, Denhardt DT, Ohnishi Y. Loss of osteopontin perturbs the epithelial-mesenchymal transition in an injured mouse lens epithelium. *Lab Invest* 2007; 87:130-8. [PMID: 17211411]
 27. Kim KK, Wei Y, Szekeres C, Kugler MC, Wolters PJ, Hill ML, Frank JA, Brumwell AN, Wheeler SE, Kreidberg JA, Chapman HA. Epithelial cell α 3 β 1 integrin links beta-catenin and Smad signaling to promote myofibroblast formation and pulmonary fibrosis. *J Clin Invest* 2009; 119:213-4. [PMID: 19104148]
 28. Lee YH, Albig AR, Regner M, Schiemann BJ, Schiemann WP. Fibulin-5 initiates epithelial-mesenchymal transition (EMT) and enhances EMT induced by TGF- β in mammary epithelial cells via a MMP-dependent mechanism. *Carcinogenesis* 2008; 29:2243-51. [PMID: 18713838]
 29. Giehl K, Menke A. Microenvironmental regulation of E-cadherin-mediated adherens junctions. *Front Biosci* 2008; 13:3975-85. [PMID: 18508491]
 30. Shintani Y, Maeda M, Chaika N, Johnson KR, Wheelock MJ. Collagen I promotes epithelial-to-mesenchymal transition in lung cancer cells via transforming growth factor- β signaling. *Am J Respir Cell Mol Biol* 2008; 38:95-104. [PMID: 17673689]
 31. Chiquet-Ehrismann R. Tenascins. *Int J Biochem Cell Biol* 2004; 36:986-90. [PMID: 15094113]
 32. Tucker RP, Chiquet-Ehrismann R. The regulation of tenascin expression by tissue microenvironments. *Biochim Biophys Acta* 2008; 1793:888-92. [PMID: 19162090]
 33. de Iongh RU, Wederell E, Lovicu FJ, McAvoy JW. Transforming growth factor- β -induced epithelial-mesenchymal transition in the lens: a model for cataract formation. *Cells Tissues Organs* 2005; 179:43-55. [PMID: 15942192]
 34. Orend G. Potential oncogenic action of tenascin-C in tumorigenesis. *Int J Biochem Cell Biol* 2005; 37:1066-83. [PMID: 15743679]
 35. Orend G, Chiquet-Ehrismann R. Tenascin-C induced signaling in cancer. *Cancer Lett* 2006; 244:143-63. [PMID: 16632194]
 36. El-Karef A, Yoshida T, Gabazza EC, Nishioka T, Inada H, Sakakura T, Imanaka-Yoshida K. Deficiency of tenascin-C attenuates liver fibrosis in immune-mediated chronic hepatitis in mice. *J Pathol* 2007; 211:86-94. [PMID: 17121418]
 37. Imanaka-Yoshida K, Matsumoto K, Hara M, Sakakura T, Yoshida T. The dynamic expression of tenascin-C and tenascin-X during early heart development in the mouse. *Differentiation* 2003; 71:291-8. [PMID: 12823230]
 38. Liu PY, Liao JK. A method for measuring Rho kinase activity in tissues and cells. *Methods Enzymol* 2008; 439:181-9. [PMID: 18374165]
 39. Jampel HD, Roche N, Stark WJ, Roberts AB. Transforming growth factor- β in human aqueous humor. *Curr Eye Res* 1990; 9:963-9. [PMID: 2276273]
 40. Cousins SW, McCabe MM, Danielpour D, Streilein JW. Identification of transforming growth factor- β as an immunosuppressive factor in aqueous humor. *Invest Ophthalmol Vis Sci* 1991; 32:2201-11. [PMID: 2071334]
 41. Saika S, Liu CY, Azhar M, Sanford LP, Doetschman T, Gendron RL, Kao CW, Kao WW. TGF β 2 in corneal morphogenesis during mouse embryonic development. *Dev Biol* 2001; 240:419-32. [PMID: 11784073]
 42. Larsson C. Protein kinase C and the regulation of the actin cytoskeleton. *Cell Signal* 2006; 18:276-84. [PMID: 16109477]
 43. Cho HJ, Yoo J. Rho activation is required for transforming growth factor- β -induced epithelial-mesenchymal

- transition in lens epithelial cells. *Cell Biol Int* 2007; 31:1225-30. [PMID: 17537651]
44. Berndt JD, Clay MR, Langenberg T, Halloran MC. Rho-kinase and myosin II affect dynamic neural crest cell behaviors during epithelial to mesenchymal transition in vivo. *Dev Biol* 2008; 324:236-44. [PMID: 18926812]
45. Matsuda A, Yoshiki A, Tagawa Y, Matsuda H, Kusakabe M. Corneal wound healing in tenascin knockout mouse. *Invest Ophthalmol Vis Sci* 1999; 40:1071-80. [PMID: 10235540]
46. Saika S, Ikeda K, Yamanaka O, Sato M, Muragaki Y, Ohnishi Y, Ooshima A, Nakajima Y, Namikawa K, Kiyama H, Flanders KC, Roberts AB. Transient adenoviral gene transfer of Smad7 prevents injury-induced epithelial-mesenchymal transition of lens epithelium in mice. *Lab Invest* 2004; 84:1259-70. [PMID: 15258599]
47. Saika S, Ikeda K, Yamanaka O, Flanders KC, Ohnishi Y, Nakajima Y, Muragaki Y, Ooshima A. Adenoviral gene transfer of BMP-7, Id2, or Id3 suppresses injury-induced epithelial-to-mesenchymal transition of lens epithelium in mice. *Am J Physiol Cell Physiol* 2006; 290:C282-9. [PMID: 16120655]