

Age-related changes in the water-soluble lens protein composition of Wistar and accelerated-senescence OXYS rats

Lyudmila V. Kopylova,¹ Ivan V. Cherepanov,^{1,2} Olga A. Snytnikova,¹ Yuliya V. Rumyantseva,³
 Nataliya G. Kolosova,³ Yuri P. Tsentlovich,¹ Renad Z. Sagdeev¹

¹International Tomography Center SB RAS, Institutskaya 3a, Novosibirsk, Russia; ²Novosibirsk State University, Pirogova 2, Novosibirsk, Russia; ³Institute of Cytology and Genetics SB RAS, Acad. Lavrentjev 10, Novosibirsk, Russia

Purpose: To determine the age-related and the cataract-specific changes in the crystallin composition in lenses of accelerated-senescence OXYS (cataract model) and Wistar (control) rats.

Methods: The water soluble (WS) and insoluble (WIS) fractions of the lens proteins were separated; the identity and relative abundance of each crystallin in WS fraction were determined with the use of two-dimensional electrophoresis (2-DE) and Matrix-Assisted Laser Desorption Ionization – Time Of Flight (MALDI-TOF) mass spectrometry. All statistical calculations were performed using the software package Statistica 6.0 by factor dispersion analysis (ANOVA/MANOVA) and Newman-Keuls post-hoc test for comparison of group mean values.

Results: The WIS protein content increased significantly in the aged animal lenses; the WIS/WS ratio increases in approximately 8 times to the age of 62 weeks. The interstrain difference was insignificant in this experiment. 2-DE maps of the young rat lenses (3 weeks) showed single spots for each lens protein while in older lenses (12 and 62 weeks) each crystallin was presented by several spots. The abundance of γ A- γ F-crystallins in WS fraction significantly decreases with age. A significant increase in the percentage abundance was also found for α -crystallins and β B2-crystallin from 3 to 12 weeks. The major differences between Wistar and OXYS lenses are the faster decay of the content of γ A- γ F-crystallins in OXYS lenses, and the significant decrease of unmodified α A-crystallin abundance in old OXYS lenses.

Conclusions: The presented results demonstrate that the increase of the water-insoluble (WIS) protein fraction is rather age-specific than cataract-specific phenomenon. The major age-related changes in WS protein composition are the fast insolubilization of γ -crystallins, and the increase of α B- and β B2-crystallin abundance. The main interstrain differences, which could be attributed to the cataract-specific processes, are the faster decay of the content of γ -crystallins and the significant decrease of unmodified α A-crystallin abundance in the OXYS lenses.

In spite of the progress made in surgical techniques in many countries during the last ten years, cataracts remain the leading cause of visual impairment globally, particularly in Asia [1]. The conditions of visual diseases are strongly age-related with earlier signs occurring in the middle age, which become more severe and more prevalent with increasing age, especially, under risk factors such as diabetes, smoking, and exposure to sunlight. The “free radical theory” of aging postulates that the major factor leading to senile cataract is the oxidation of the lens tissues by reactive oxygen species (ROS) [2].

Lens is a unique organ because of its composition and properties. Lens proteins constitute more than 30% of the lens total mass [3-5]. The major constituents of the lens are crystallins – the structural proteins that amount to about 90% of the total proteins in the lens [6,7]. There are three types of crystallins in the mammalian lens – α -, β -, and γ -crystallins, with a high level of homology within each group [8]. The transparency of the eye lens strongly depends on the crystallin

solubility and structure [9-11]. All crystallins are initially water-soluble (WS). During aging, crystallins may undergo post-translational modifications (PTMs) which accumulate throughout life, since the turnover of crystallins in the lens is negligible [12]. Modifications of the lens crystallins disrupt its normal structure in the lens: proteins undergo oxidation, coloration, aggregation, and lose the solubility. Crystallins start forming large insoluble aggregates that may exceed the dimension of the wavelength of light and produce fluctuations in the refraction index causing light scattering [3,5,13,14]. These processes eventually result in the lens opacification and cause pathological alterations leading to the cataract formation [15]. The well known PTMs are the protein truncation, deamidation, glycation, disulfide formation, cross-linking, and phosphorylation [6,7,10,11,16,17]. The most significant influence on the protein insolubility is probably caused by the PTMs of α -crystallins: in the tissues α -crystallins appear to have chaperone activity which may decrease due to modifications [8,18-21]. Other crystallins undergo similar changes, and all these processes lead to the increasing percent of water-insoluble (WIS) fraction during aging [13,22,23]. In several studies, it was shown that the listed modifications are common for primates and rodents

Correspondence to: Yuri P. Tsentlovich, International Tomography Center, 630090, Institutskaya 3a, Novosibirsk, Russia; Phone: +7-383-3303136; FAX: +7-383-3331399; email: yura@tomo.nsc.ru

such as mice [24-26] and rats [27-29]. At present, it is not completely clear which modifications are cataract-specific, and which are just a part of the normal maturation and aging processes (such as site-specific proteolysis of β -crystallins in young lenses for tighter packing of crystallins during maturation [17,27]). It is necessary to have both normal and cataractous lenses of different ages to compare the obtained results and to be able to detect disease-specific modifications. Obviously, normal human lenses are difficult to obtain, especially from the young donors. Thus, different rodent models have been developed to study the aging processes [30-34]. None of these models exactly matches to what occurs in the human lens, but the mechanisms of cataract formation are supposed to be similar.

In this study, we used the strain of senescence-accelerated OXYS rats, which appears to be a good model of senile cataract [35-37]. This strain was developed at the Institute of Cytology and Genetics SB RAS, Novosibirsk, Russia, from Wistar stock by selection for their susceptibility to the cataractogenic effect of galactose [38]. It was shown that the first signs of cataract in OXYS rats appear at the age of 1.5 months, to the age of 3 months 90% of animals are affected by the lens opacification, and at the age of 4-6 months the morbidity reaches 100% [36,39]. The mature cataract was detected in 90% of eyes of two-year-old OXYS rats, whereas the same aged Wistar rats showed only the initial signs of cataract [39].

The purpose of the present study was to analyze the water-soluble protein content of OXYS and Wistar rat lenses and to determine its age-dependent alterations. In this work, we obtained 2-DE protein maps for lenses of increasing age and of two rat strains to serve as reference for further investigations. The protein identities were confirmed by mass-spectrometry (MALDI) and each protein relative content was quantified. The obtained data will facilitate the future experiments on detecting the cataract-specific modifications in rat lenses.

METHODS

Materials and reagents: Phosphate buffer tablets (Biolot, Saint-Petersburg, Russia), urea (Bio-Rad Laboratories, Hercules, CA), acrylamide (4K; Medigen Laboratories, Novosibirsk, Russia), acrylamide for IEF (Amersham Biosciences, Uppsala, Sweden), bis-acrylamide (Amresco, Solon, OH), ampholites (Bio-Lyte 3/10; Bio-Lyte 5/8; Bio-Rad), CHAPS (Bio-Rad), ammonium persulphate (PSA; Helicon, Moscow, Russia), TEMED (Helicon), NaOH (Reachim, Moscow, Russia), orthophosphoric acid (Reachim), glycerol (Panreac, Barcelona, Spain), Tris-HCl (Bio-Rad), glycine (Bio-Rad), SDS (Bio-Rad), DTT (Helicon), bromophenol blue (Helicon), agarose (Bio-Rad), Coomassie brilliant blue R-250 (Sigma, Steinheim, Germany), acetic acid (Chimreactiv, Moscow, Russia), ammonium bicarbonate (AMB; Fluka, Steinheim, Germany),

ACN (Cryochrom, Saint-Petersburg, Russia), sequencing grade modified trypsin (Promega, Madison, WI), TFA (Sigma), 2,5-DHB (Bruker Daltonics, Bremen, Germany) were used as received. H₂O was deionized.

Rat lens preparation: All animal procedures adhered to the Association for Research in Vision and Ophthalmology (ARVO) statement for the Use of Animals in Ophthalmic and Vision Research and in compliance with the European Communities Council Directive No. 86/609/EES. Lenses were obtained from Wistar and senescent-accelerated OXYS rat strains maintained in the Breeding Laboratory of the Institute of Cytology and Genetics SB RAS, Novosibirsk, Russia. Animals were put to sleep with use of diethyl ester and decapitated. The lenses were removed, frozen in liquid nitrogen, and stored at -70 °C until analysis. The content of the water-soluble and water-insoluble lens proteins was determined using lenses from Wistar and OXYS rats at the age of 3, 12, 54, and 62 weeks. Lenses were homogenized in 600 μ l of 0.02 M phosphate buffer solution, pH 7.3, containing protease inhibitor (Protease Inhibitor Cocktail; Sigma). The WS and WIS fractions were separated by centrifugation at 12,000 \times g for 50 min at 4 °C. After separation, the supernatant and the pellet were dried in a vacuum evaporator (Christ AVC 2-25 CD Plus; 3000 g; 37 °C; Martin Christ Gefriertruhenanlagen GmbH, Osterode am Harz, Germany), and then weighted. Clean buffer solution was also dried and weighted, and the mass of the buffer salts was subtracted from the masses of WS and WIS protein fractions.

The WS proteins for electrophoresis were extracted from 36 lenses from Wistar and OXYS rats at the age of 3, 12, and 62 weeks. The number of animals in each group varied from 3 to 6. The proteins were isolated from lenses by homogenization on ice in 0.02 M phosphate buffer solution, pH 7.3, containing protease inhibitor (Protease Inhibitor Cocktail; Sigma). The volume of the buffer solution for 12- and 62-week-old rats was 600 μ l per lens (only one lens from each animal was used). Since 3-week-old lenses are much smaller than the mature ones, the homogenization was performed using both lenses from each animal of this age, and the buffer solution volume was 300 μ l per lens pair. The WS and WIS fractions were separated by centrifugation at 12,000 \times g for 50 min at 4 °C. The protein content in supernatants was determined using Bradford reagent [40] (Fermentas Inc., Glen Burnie, MD) and BSA standard (Bio-Rad) following the manufacturer's protocol.

RNA isolation, reverse transcription, preparation of "standard" cDNA and real time PCR were described in details earlier [39]. Briefly, total cell RNA was isolated from the rat lens by the phenol-chloroform method. Expression of alpha-crystallin A chain (*Cryaa*) and alpha-crystallin B chain (*Cryab*) genes was determined by real time PCR, the housekeeping gene 60S ribosomal protein L30 (*Rpl30*) was used as a reference gene.

TABLE 1. CONTENT OF WS AND WIS PROTEINS IN LENSES OF WISTAR AND OXYS RATS.

Age	Strain	Lens weight, mg	WS proteins mass, mg	WIS proteins mass, mg	Ratio WIS/WS
3 weeks	Wistar	16.0±0.7	5.8±0.2	0.65±0.13	0.11
	OXYS	17.8±0.9	6.3±0.2	0.64±0.15	0.10
12 weeks	Wistar	37.6±1.9	12.9±1.6	4.9±0.2	0.38
	OXYS	41.7±2.1	13.9±0.6	4.3±0.4	0.31
54 weeks	Wistar	69.3±4.3	21.4±1.6	12.7±1.6	0.60
	OXYS	68.5±2.4	21.4±1.0	11.4±0.5	0.53
62 weeks	Wistar	79.4±4.4	18.6±1.1	15.9±1.5	0.85
	OXYS	64.3±5.2	16.3±1.7	12.9±0.9	0.79

Two-dimensional gel electrophoresis: Isoelectric focusing (IEF) was performed using “tube gel” system. The gel for the first dimension contained 1.75 g urea, 1,190 µl H₂O, 455 µl of stock solution containing 30% acrylamide for IEF and 0.8% bis-acrylamide, 62 µl ampholites 3/10, 123 µl ampholites 5/8, 210 µl 30% CHAPS, 7 µl 10% PSA (freshly prepared), and 3.5 µl TEMED. The final percentage of acrylamide in tube gels was approximately 4%. PSA and TEMED were added to the solution directly prior casting the gel. The gels were cast in glass tubes (length 20 cm, internal diameter 1.5 mm). The length of gels in tubes was 18 cm. After the gel polymerization, 150 µg (in 10 µl) of WS protein mixture (prepared as written above) from each sample was loaded onto the top of the gels followed by 0.05 M NaOH (~10 µl) to fill the tube up. The bottom of the tube was covered with a drop of 0.13% orthophosphoric acid to dislodge any bubbles from inside the gel tubes. The tube stand with the tubes was fixed on a cooling core and placed into the lower tank of a Protean II xi 2-D Cell (Bio-Rad) filled with 0.13% orthophosphoric acid. The upper tank was filled with 400 ml of 0.05 M NaOH. Running conditions for IEF were: 100 V to 600 V over 1 h, 700 V over 10 h, 900 V over 1 h, room temperature.

After the IEF stage, the gels were extruded from the tubes into the tray with buffer solution containing 6 M urea, 30% glycerol, 12.5 mM Tris-HCl (pH 6.8), 2% SDS, a trace of bromophenol blue; 0.2% DTT was added immediately before use. The gels were left in the solution for 5–10 min, and then placed over the 12% SDS-PAGE gel for the second dimension. The gels were covered with 0.9% agarose containing a trace of bromophenol blue tracking dye. Slabs for the second dimension were prepared in advance. SDS-PAGE gels (12%; 20×20 cm, thickness 1.5 mm) contained 120 ml of stock solution (30% acrylamide and 0.8% bis-acrylamide), 1.5 M Tris-HCl with 0.4% SDS (76 ml), 10% PSA (682 µl), TEMED (123 µl) and H₂O (103 ml). The prepared slabs were fixed on a cooling core and placed into the lower tank of a Protean II Multi-Cell (Bio-Rad) filled with electrophoresis buffer (25 mM Tris (pH 8.3), 192 mM glycine, 0.1% SDS). Running conditions for the second dimension

were: 20 mA per gel (for 20 min), 40 mA per gel (for 2 h), 35 mA per gel (for 3 h). After electrophoresis, the gels were stained with 0.025% Coomassie blue R-250 in 10% acetic acid.

Protein quantification and identification: Gel images were obtained using a VersaDoc Imaging System (4000 MP; Bio-Rad), and calculation of the protein percentage abundances was performed with a PDQuest Advanced 2D-analysis Software 8.0.1 (Bio-Rad). Proteins from gels were identified by MS analysis. Coomassie stained spots from gels (for each strain and each age of animals) were manually excised (including negative control, no protein) and washed twice in 0.2 M AMB buffer in 50% ACN at 37 °C to remove the stain. Each gel piece was dried with 100% ACN and digested with sequencing grade modified trypsin (12.5 ng/µl) plus 40 mM AMB for 16 h at 37 °C. Peptides were desalted using C18 ZipTips (Millipore Corporation, Billerica, MA), mixed (in a ratio of 0.5 µl of the sample to 0.5 µl of the matrix) with 2.5-DHB dissolved in 70% ACN/0.1% TFA and spotted to a standard MTP ground steel plate (Bruker Daltonics, Bremen, Germany). The further analysis was performed using a MALDI-TOF/TOF spectrometer Ultraflex III (Bruker Daltonics). The mass spectra of protein tryptic digests were recorded in reflective positive ion mode in the 500–4200 m/z range. Spectra were then analyzed using a FlexAnalysis software 3.0 (Build 96; Bruker Daltonics), and peptide masses were entered into the local MASCOT server 2.2.04 for the identification of peptides. The MALDI-TOF identities of proteins were established by using the SwissProt database (updated 2010–08–10; mass accuracy – 70 ppm, 1 missed cleavage).

Statistical analysis: All statistical calculations were made using the software package Statistica 6.0 (Statsoft Russia, Moscow, Russia) using factor dispersion analysis (ANOVA/MANOVA) and Newman-Keuls post-hoc test for comparison of group mean values. Genotype and age of animals were considered as independent factors. In all cases, results were considered as statistically significant at p<0.05.




Figure 1. 2-DE gel of the lens proteins of 12-week-old Wistar rat stained with Coomassie brilliant blue R-250. Arrows indicate the unmodified proteins, and their assignment is shown, the modified crystallins are given the numbers only: spots 2–5,7,8 correspond to α A-crystallin, spots 10,11 – to β A4-crystallin, spots 15–20 – to β A3-crystallin, spot 22 – to β B1-crystallin, spot 23 – to β B3-crystallin, spots 26–27 – to β B2-crystallin, spot 28 – to α B-crystallin, spot 31 – to γ E-crystallin and spot 33 – to γ D-crystallin. The assignment of all indicated spots and the sequence coverage are also presented in Table 2.

RESULTS

Age-related changes in WIS/WS protein ratio: The measurements of the WS and WIS protein content in the lens of OXYS and Wistar rats of different age were performed for lenses of three animals for every age and every strain. The analysis was performed for every lens separately, the averaged results are presented in Table 1. For both strains, a significant increase of the WIS protein content is observed: the WIS/WS ratio increases from approximately 0.1 (3 weeks), to 0.4 (12 weeks), to 0.8 (62 weeks). The interstrain difference for all ages is insignificant: in fact, for all ages the WIS/WS ratio for Wistar rats is slightly higher than that for senescence-accelerated OXYS rats, but this difference is below the experimental error.

2-DE maps of water-soluble proteins: To determine the relative abundance, the identities of crystallin subunits, and the age-related differences between OXYS and Wistar rat strains, 2-DE was performed, the spots were visualized and identified by in-gel digestion with the subsequent MS analysis. Figure 1 shows a 2-DE map of the water-soluble protein fraction from the lens of a 12-week-old Wistar rat, the assignments of the major proteins are indicated. The identities of proteins were determined with 60%–80% of sequence coverage by MALDI-MS analysis of tryptic peptides. The results obtained for the WS fraction of 12-week-old Wistar lens are given in Table 2, the same procedure was applied to proteins of all ages and both strains. In 2-DE maps of 3-week-old lenses, most of the crystallins are represented by a single spot, while in older lenses the number of spots significantly increases. Thus, the spots present in the maps of young lenses are attributed to the unmodified proteins, while the new spots observed in the maps of older lenses correspond to the crystallins, which underwent post-translational

modifications. In Figure 1, arrows indicate the unmodified proteins, and their assignment is shown, whereas the modified crystallins are given the numbers only: spots 2–5, 7, and 8 correspond to modified α A-crystallins, spots 10–11 – to modified β A4-crystallins, spots 15–20 – to modified β A3-crystallins, spot 22 – to modified β B1-crystallin, spot 23 – to modified β B3-crystallin, spots 26–27 – to modified β B2-crystallins, spot 28 – to modified α B-crystallin, spot 31 – to modified γ E-crystallin and spot 33 – to modified γ D-crystallin. The obtained 2-DE maps are in a good agreement with the crystallin maps of rats and mice [25,29] published earlier. In addition, we were able to obtain a good resolution for γ -crystallins.

Two non-crystallin proteins were also identified in the soluble fraction of the lens proteins. One of them is identified as grifin (molecular weight 15,839 Da, accession number [O88644](#) (GRIFN_RAT) at [Uniprot](#). This galectin-related inter-fiber protein is lens-specific, it is located at the interface between lens fiber cells. This protein has been previously described in rat lenses and has a molecular function of sugar binding [41]. In our samples, its level (0.4%–1% of the total protein content) is comparable with the levels of the least abundant crystallins. The other protein found in the rat lens is fatty acid-binding protein (molecular weight 15,059 Da, accession number [P55053](#) (FABP5_RAT) at [Uniprot](#). This protein plays a role of the marker for differentiation of the fiber cells. It was previously found in the mice [25], cow [42], and rat [43] lenses.

Age-related changes in the crystallin composition: Age-related changes in the protein composition of rat lenses were monitored using the water-soluble lens proteins from 3-, 12-, and 62-week-old Wistar and OXYS rats. For each age, the lenses from 3–6 animals were taken, the water-soluble protein

TABLE 2. IDENTIFIED PROTEINS FROM THE WS FRACTION OF 12-WEEK-OLD WISTAR LENS WITH SEQUENCE COVERAGE (%) AFTER MASCOT SEARCH.

Spot number	Identified protein	Sequence coverage, %	Spot number	Identified protein	Sequence coverage, %
1	Grifin	28	21	βB1	63
2	α A	53	22	β B1	65
3	α A	64	23	β B3	75
4	α A	66	24	βB3	70
5	α A	53	25	βB2	50
6	αA	90	26	β B2	72
7	α A	86	27	β B2	69
8	α A	58	28	α B	42
9	αA_{insert}	39	29	αB	67
10	β A4	69	30	γS	74
11	β A4	63	31	γ E	63
12	βA4	63	32	γE	73
13	βA2	56	33	γ D	67
14	βA3	62	34	γD	54
15	β A3	48	35	γF	76
16	β A3	51	36	γC	76
17	β A3	51	37	γB	94
18	β A3	69	38	γA	61
19	β A3	67	39	FABP	70
20	β A3	62			

Unmodified proteins are indicated by bold fonts.

fraction from each lens was subjected to 2-DE (Figure 2), and the percentage abundance of crystallins observed in gels was determined by the integral density of the corresponding spots. The data obtained for the same age and strain were averaged. Since the spots corresponding to γ A- and γ B-crystallin in some gels overlap, the percentage abundances of these crystallins were combined. The average percentage abundance of unmodified proteins are given in Table 3 and Figure 3, and the data on the sum of modified and unmodified crystallins are collected in Table 4 and presented in Figure 4.

The most dramatic and rapid age-related changes were registered for γ -crystallins: a fast decay of the concentration is observed for all γ -crystallins (except γ S-crystallin). The decay of γ -crystallin in lenses of OXYS rats is faster than that in Wistar. The total content of γ A- γ F-crystallin in Wistar decays from 61.4% (3 weeks) to 40.6% (12 weeks) to 9.5% (62 weeks), while in OXYS lenses the age dynamics of γ A- γ F-crystallin is: 65.9% (3 weeks) – 28% (12 weeks) – 7% (62 weeks). The fastest decay is observed for γ F-crystallin, which level decreases from 10.8% (3 weeks) to 1.3% (12 weeks) in Wistar and from 10.4% (3 weeks) to 0.3% (12 weeks) in OXYS lenses. At the age of 62 weeks, γ F-crystallin have not been detected neither in Wistar nor in OXYS lenses. It is important to note that in all gels the majority of γ -crystallin is

present in unmodified state. That likely means that the modifications of γ -crystallins result in their fast insolubilization.

Against a background of the strong γ -crystallin decay, the percentage abundance of other crystallins (with the few exceptions) either increases with age, or remains approximately at the same level. The most pronounced increase is observed for α B- and β B2-crystallin: the percentage of these proteins in 62-week-old lenses is higher than in 3-week-old lenses by approximately an order of magnitude. This effect should be attributed to the enhanced stability of these proteins under oxidative stress. However, the increased expression of α B- and β B2-crystallin in the epithelial cells of the lens also cannot be ruled out. The difference between α B- protein contents in the Wistar and OXYS lenses at 3 weeks and 12 weeks is insignificant, but at 62 weeks the percentage abundance of α B-crystallin in Wistar is twice as large as in OXYS lenses, mostly due to the high abundance of modified α B-crystallin in the WS fraction of proteins from the Wistar lenses. The most plausible explanation of this observation is that at the age of 62 weeks, the modifications of α B-crystallin in OXYS lenses become too severe, and they undergo insolubilization.




Figure 2. 2-DE gels of WS fractions of the lens proteins of 3, 12, and 62-week-old Wistar and OXYS rats. Assignment is given for unmodified crystallins.

An interesting dynamics is observed for α A-crystallin. During the period from 3 weeks to 12 weeks, the sum of modified and unmodified α A-crystallin in lenses of Wistar rats increases from 9.6% to 20.2%, and in OXYS from 6.7% to 28.1%. At 62 weeks, the percentage of α A-crystallin in Wistar remains approximately the same as in 12 weeks, while in OXYS lenses it decreases to 18.4%. At this age, unmodified α A-crystallin constitutes about 60% of the total amount of α A-crystallin in Wistar lenses, and only approximately 45% in OXYS.

DISCUSSION

2-DE maps of young rat lenses obtained in the present work are in a good agreement with the previously reported proteome maps of lenses of Sprague-Dawley rats [29] and FVB/N mice [25]. The position, masses and pIs of crystallins from rats of different strains and from mice are very similar, which indicates the absence of differences in the initial steps of rodent's lens development. The general content of WS proteins in young lenses from Wistar and Sprague-Dawley rats is also similar: approximately 10% of α -crystallin, 30% of β -crystallin, and 60% of γ -crystallin [29]. Nevertheless, the percentage abundances of some individual proteins are

different. The most significant difference was found for α B-crystallin, which content in 3-week-old Wistar and OXYS lenses (0.5%–0.8%) is much lower than that in 16-day-old Sprague-Dawley lens (approximately 4% [29]).

The time evolution of the lens protein composition includes the changes in the contribution of α -, β -, and γ -crystallin families into total protein content as well as in the abundance of individual crystallins. The major factor, which influences the protein composition in WS fraction, is the insolubilization of γ -crystallin. This process occurs in both Wistar and OXYS lenses (although in OXYS it proceeds faster) and results in the decrease of the γ A- γ F-crystallin percentage abundance down to 7%–10% to the age of 62 weeks. γ -Crystallins are fully presented in rodent lenses, and their intensive degradation during aging has been reported for both mice [25] and rats [44]. In human lenses, there are three types of γ -crystallin: γ C-, γ D-, and γ S-crystallin [45]. It has been shown [17] that the concentration of γ C- and γ D-crystallin in human lens also decreases with age. The major process causing the γ -crystallin insolubilization might be the extensive deamidation of crystallins, which enhances the acidity of proteins [18], and the truncation [46].

TABLE 3. PERCENTAGE ABUNDANCE OF UNMODIFIED PROTEINS IN LENSES OF WISTAR AND OXYS RATS.

Age	3 weeks		12 weeks		62 weeks	
	Wistar	OXYS	Wistar	OXYS	Wistar	OXYS
Strain						
Number of animals	5	6	4	3	4	3
α A	9.6 \pm 0.9	6.7 \pm 1.8	13.8 \pm 3.0	19.5 \pm 1.0	13.2 \pm 1.5	8.5 \pm 0.9
α B	0.5 \pm 0.3	0.8 \pm 0.6	1.8 \pm 0.3	2.3 \pm 0.7	4.5 \pm 0.6	5.9 \pm 0.7
β A2	1.2 \pm 0.7	0.4 \pm 0.4	3.2 \pm 0.4	4.3 \pm 2.1	0	0
β A3	3.0 \pm 0.5	1.6 \pm 1.6	1.2 \pm 0.2	0.7 \pm 0.1	0.9 \pm 0.1	2.4 \pm 1.8
β A4	3.3 \pm 1.0	3.2 \pm 1.3	3.9 \pm 0.7	3.7 \pm 0.6	4.6 \pm 1.9	3.6 \pm 1.2
β B1	4.3 \pm 0.9	3.7 \pm 0.8	1.1 \pm 1.1	2.6 \pm 0.4	0.9 \pm 0.4	0.8 \pm 0.5
β B2	3.5 \pm 0.5	3.4 \pm 1.3	9.4 \pm 2.6	15.0 \pm 5.8	13.4 \pm 3.1	19.1 \pm 0.8
β B3	5.0 \pm 0.6	4.5 \pm 1.3	0	0	1.2 \pm 0.4	1.3 \pm 0.1
γ A + γ B	17.8 \pm 2.6	24.5 \pm 6.6	14.7 \pm 5.4	9.1 \pm 5.9	4.9 \pm 1.6	2.5 \pm 1.4
γ C	6.7 \pm 2.1	4.2 \pm 1.7	8.0 \pm 4.2	2.9 \pm 4.4	2.3 \pm 1.3	1.4 \pm 1.1
γ D	8.0 \pm 1.4	8.4 \pm 1.1	9.1 \pm 2.2	7.7 \pm 3.9	1.4 \pm 0.4	0.8 \pm 0.5
γ E	7.4 \pm 2.0	8.9 \pm 2.2	4.3 \pm 1.9	2.4 \pm 2.2	0	0.7 \pm 0.1
γ F	7.5 \pm 1.6	7.5 \pm 2.7	0	0.9 \pm 0.1	0	0
γ S	2.1 \pm 0.7	1.6 \pm 0.4	5.1 \pm 1.9	5.1 \pm 3.7	4.4 \pm 1.1	9.8 \pm 5.3

TABLE 4. PERCENTAGE ABUNDANCE OF THE SUM OF MODIFIED AND UNMODIFIED PROTEINS IN LENSES OF WISTAR AND OXYS RATS.

Age	3 weeks		12 weeks		62 weeks	
	Wistar	OXYS	Wistar	OXYS	Wistar	OXYS
Strain						
Number of animals	5	6	4	3	4	3
α A	9.6 \pm 0.9	6.7 \pm 1.8	20.2 \pm 3.4	28.1 \pm 1.4	21.3 \pm 1.3	18.4 \pm 2.7
α B	0.5 \pm 0.3	0.8 \pm 0.7	3.1 \pm 1.0	4.3 \pm 2.2	11.3 \pm 3.0	6.6 \pm 0.8
β A2	1.2 \pm 0.7	0.4 \pm 0.4	2.4 \pm 1.6	4.3 \pm 2.1	0	0
β A3	12.8 \pm 0.7	8.3 \pm 3.6	8.4 \pm 4.0	5.9 \pm 0.7	7.5 \pm 0.9	8.5 \pm 2.2
β A4	3.3 \pm 1.0	3.2 \pm 1.5	4.9 \pm 1.5	4.5 \pm 0.4	11.0 \pm 3.7	9.0 \pm 5.1
β B1	4.7 \pm 1.1	3.8 \pm 0.7	1.3 \pm 1.6	2.6 \pm 0.4	2.7 \pm 1.3	5.3 \pm 4.2
β B2	3.5 \pm 0.5	3.4 \pm 1.3	11.5 \pm 3.2	19.0 \pm 5.2	24.5 \pm 6.3	29.5 \pm 4.4
β B3	5.5 \pm 0.4	5.8 \pm 1.2	3.6 \pm 0.7	2.1 \pm 1.2	2.8 \pm 0.8	3.8 \pm 0.8
γ A + γ B	19.0 \pm 1.8	24.8 \pm 3.2	14.7 \pm 6.1	9.1 \pm 5.9	4.9 \pm 1.6	3.2 \pm 2.1
γ C	6.7 \pm 2.1	4.2 \pm 1.7	8.0 \pm 4.2	2.9 \pm 4.4	2.8 \pm 1.0	1.8 \pm 1.6
γ D	14.4 \pm 2.6	14.1 \pm 3.7	13.0 \pm 2.5	12.7 \pm 3.8	1.4 \pm 0.4	1.2 \pm 1.0
γ E	10.4 \pm 2.0	12.4 \pm 1.4	5.5 \pm 2.2	3.0 \pm 2.0	0.5 \pm 0.2	0.7 \pm 0.6
γ F	10.8 \pm 1.5	10.4 \pm 3.3	1.3 \pm 0.9	0.3 \pm 0.5	0	0
γ S	2.1 \pm 0.7	1.6 \pm 0.4	5.1 \pm 1.9	5.1 \pm 3.7	4.4 \pm 1.1	9.8 \pm 5.3
others	0.1 \pm 0.2	2.3 \pm 1.8	1.1 \pm 0.8	1.9 \pm 2.0	4.8 \pm 1.6	2.2 \pm 1.2

Very likely that the insolubilization of γ -crystallin is the main cause of the age-related changes in WIS/WS ratio (Table 1): at the age of 3 weeks, WS proteins constitute 30%–40% of the total lens weight, and WIS proteins – only 3%–5%; at the age of 62 weeks, the contribution of WS fraction decreases to approximately 25%, and that of WIS fraction increases to about 20%. The age-related changes in WIS/WS ratio in the Wistar lens has been recently reported [5]: from the age of 16 days to 1 year, this ratio increases by approximately a factor of four, which is in a good agreement with our data. We should note that the absolute values of WIS/WS ratio in work [5] were significantly lower than in the present work – probably, due to the different procedures of the WS and WIS fraction measurement.

The insolubilization of the lens protein is often referred to as one of the main factors of the cataract development: insoluble proteins may form large aggregates causing light scattering and lens opacification. Our results suggest that the increase of the WIS fraction is the age-specific rather than cataract-specific process: the WIS/WS ratio in Wistar and senescence-accelerated OXYS rat lenses is approximately the same for all ages, though the onset of the cataract in OXYS rats occurs much earlier [39].

From 3 weeks to 12 weeks, the most significant increase in the percentage abundance was found for both α -crystallin and β B2-crystallin. In the lens, α A-crystallin plays a very important role due to its chaperone activity [8,18,47]: with aging, α -crystallin binds partially unfolded β - and γ -

crystallins, thereby preventing their further aggregation and precipitation, and delaying lens opacification [9,47]. Thus, the increase of the α -crystallin content can be attributed to the compensatory reaction of the lens to the oxidative stress, which causes the enhanced expression of α -crystallin. To confirm this assumption, we performed the comparative measurement of the α -crystallin expression in young (3-week-old) Wistar and OXYS lenses using real time PCR. The expression of α -crystallin was measured for lenses of 7 Wistar and 7 OXYS rats using the housekeeping gene *Rpl30* as a reference gene (see Experimental section), the results were averaged. It has been found that the expression of both α A- and α B-crystallin ($E_{\alpha A}$ and $E_{\alpha B}$) in OXYS lenses is twice as large as that in Wistar lenses: $E_{\alpha A}(\text{Wistar})=2.45\pm 0.28$; $E_{\alpha A}(\text{OXYS})=4.87\pm 0.27$; $E_{\alpha B}(\text{Wistar})=3.04\pm 0.21$;

$E_{\alpha B}(\text{OXYS})=6.11\pm 0.53$. Since tissues in OXYS rats are more exposed to the oxidative stress than ones in Wistar rats, the compensatory reactions should be stronger. Thus, the more significant growth of the α -crystallin content in OXYS lenses in comparison with that in Wistar rats should be attributed to the enhanced expression of α -crystallin as a compensatory reaction to the oxidative stress. PTMs cause breaches in the protein folding, and accumulation of proteins with the “wrong” folding can initiate unfolded protein response, which implies an accelerated degradation of “wrong” proteins as well as a deceleration of protein synthesis [48]. This mechanism can explain age-related decrease in synthesis of some proteins.

The content of α A-crystallin in Wistar lenses at 62 weeks is approximately the same as at 12 weeks, whereas in OXYS




Figure 3. Relative abundances of unmodified soluble crystallins in lenses of Wistar and OXYS rats at the ages of 3, 12 and 62 weeks. Black bars show the values for 3-week-old animals, hatched bars – for 12-week-old animals, and open bars – for 62-week-old animals. Error bars indicate the standard deviation.




Figure 4. Relative abundances of total soluble proteins in lenses of Wistar and OXYS rats at the ages of 3, 12 and 62 weeks. Black bars show the values for 3-week-old animals, hatched bars – for 12-week-old animals, and open bars – for 62-week-old animals. Error bars indicate the standard deviation.

lenses it decreases with age. This decrease is especially pronounced for unmodified α A-crystallin, which percentage abundance in 62-week-old OXYS lenses is less than a half from that in Wistar lenses of the same age. This finding correlates with the recent report on very low α A-crystallin expression in old OXYS lenses [39], which likely can be attributed to the damages affecting the epithelial cell functioning. Another possible explanation of the faster α A-crystallin loss during aging than that of α B-crystallin is that α A-crystallin is more vulnerable to the oxidative stress. This hypothesis is confirmed by the recent studies of α -crystallin distribution in the bovine lenses. It has been shown [7] that α B-crystallin is stable on both the periphery and the nucleus of the bovine lens, which means that this protein should accumulate with age. On the contrary, intact α A-crystallin was found only on the periphery of bovine lens, where new proteins are synthesized [18,47]. The three major truncated forms α A₁₋₁₀₁, α A₁₋₆₅, α A₁₋₅₈ were observed across the whole section and especially in the nucleus where older proteins accumulate with age [7].

The content of β B2-crystallin in WS fraction of the lens proteins undergoes a stable growth throughout the lifespan of both Wistar and OXYS rats, demonstrating the high stability of this protein under oxidative stress. It was suggested that with the loss of α A-crystallin, it might be substituted in the lens structure by β B2-crystallin [45].

In summary, the results of this work show that the most pronounced age-related changes in the protein composition of the rat lens are the increase of WIS/WS ratio with aging, the fast insolubilization of γ A- γ F-crystallins, and the increase of α B- and β B2-crystallin abundance in the WS protein fraction during the lens growth. The percentage abundance of α A-crystallin in WS fraction grows only in relatively young lenses. The major differences between Wistar and OXYS lenses are the faster decay of the content of γ A- γ F-crystallin in OXYS lenses, and the significant decrease of unmodified α A-crystallin abundance in old OXYS lenses. These differences can be attributed to the cataract-specific changes in the protein composition of the lens. More detailed information on the cataract-specific processes in the lens can be obtained from the analysis of the content of the WIS fraction of lens proteins, and from the comparative study of post-translational modifications of proteins in Wistar and OXYS lenses. This work is currently in progress in our laboratory.

ACKNOWLEDGMENTS

This work was supported by the following agencies: FASI state contracts 14.740.11.0758, 02.740.11.0262 and P708, Russian Foundation for Basic Research (projects 11-04-00143-a and 11-03-00296-a), President of the Russian Federation (grant NSh-7643.2010.3), the Division of Chemistry and Material Science, Russian Academy of Sciences, and the Government of Moscow (Program “New

Methods and Technologies of Early Diagnostics and treatment of Oncological and Other Dangerous Diseases”).

REFERENCES

1. Foster A, Gilbert C, Johnson G. Changing patterns in global blindness: 1998–2008. *Community Eye Health* 2008; 21:37-9. [PMID: 19030126]
2. Fletcher AE. Free radicals, antioxidants and eye diseases: evidence from epidemiological studies on cataract and age-related macular degeneration. *Ophthalmic Res* 2010; 44:191-8. [PMID: 20829643]
3. Benedek GB. Theory of transparency of the eye. *Appl Opt* 1971; 10:459-73. [PMID: 20094474]
4. Graw J. The crystallins: genes, proteins and diseases. *Biol Chem* 1997; 378:1331-48. [PMID: 9426193]
5. Sakthivel M, Elanchezian R, Thomas PA, Geraldine P. Alterations in lenticular proteins during ageing and selenite-induced cataractogenesis in Wistar rats. *Mol Vis* 2010; 16:445-53. [PMID: 20300567]
6. Groenen PJ, Merck KB, de Jong WW, Bloemendal H. Structure and modifications of the junior chaperone α -crystallin. *Eur J Biochem* 1994; 225:1-19. [PMID: 7925426]
7. Han J, Schey K. MALDI tissue imaging of ocular lens α -crystallin. *Invest Ophthalmol Vis Sci* 2006; 47:2990-6. [PMID: 16799044]
8. Bloemendal H, de Jong W, Jaenicke R, Lubsen NH, Slingsby C, Tardieu A. Ageing and vision: structure, stability and function of lens crystallins. *Prog Biophys Mol Biol* 2004; 86:407-85. [PMID: 15302206]
9. Horwitz J. Alpha-crystallin can function as a molecule chaperone. *Proc Natl Acad Sci USA* 1992; 89:10449-53. [PMID: 1438232]
10. Horwitz J. Alpha-crystallin. *Exp Eye Res* 2003; 76:145-53. [PMID: 12565801]
11. Reddy GB, Das KP, Petrash JM, Surewicz WK. Temperature-dependent chaperone activity and structural properties of human alphaA- and alphaB-crystallins. *J Biol Chem* 2000; 275:4565-70. [PMID: 10671481]
12. Young RW, Fulhorst HW. Regional differences in protein synthesis within the lens of the rat. *Invest Ophthalmol* 1966; 5:288-97. [PMID: 5947946]
13. Spector A. The search of a solution to senile cataract. *Invest Ophthalmol Vis Sci* 1984; 25:130-46. [PMID: 6321383]
14. Clark JI. Development and maintenance of lens transparency. In: Albert DA, Jacobiec FA, editors. *Principles and practice of ophthalmology*. Philadelphia: WB Saunders Company; 1994. p. 114–23.
15. Schachar RA, Solin SA. The microscopic protein structure of the lens with a theory for cataract formation as determined by Raman spectroscopy of intact bovine lenses. *Invest Ophthalmol* 1975; 14:380-96. [PMID: 1126827]
16. Miesbauer LR, Zhou X, Yang Z, Yang Z, Sun Y, Smith DL, Smith JB. Post-translational modifications of water-soluble human lens crystallins from young adults. *J Biol Chem* 1994; 269:12494-502. [PMID: 8175657]
17. Lampi KJ, Ma Z, Hanson SR, Azuma M, Shih M, Shearer TR, Smith DL, Smith JB, David LL. Age-related changes in human lens crystallins identified by two-dimensional

- electrophoresis and mass spectrometry. *Exp Eye Res* 1998; 67:31-43. [PMID: 9702176]
18. Sharma KK, Santhoshkumar P. Lens aging: Effects of crystallins. *Biochim Biophys Acta* 2009; 1790:1095-108. [PMID: 19463898]
 19. Andley UP, Mathur S, Griest TA, Petrash JM. Cloning, expression, and chaperone-like activity of human alpha A-crystallin. *J Biol Chem* 1996; 271:31973-80. [PMID: 8943244]
 20. Bova MP, Ding LL, Horwitz J, Fung BKK. Subunit exchange of alpha A-crystallin. *J Biol Chem* 1997; 272:29511-7. [PMID: 9368012]
 21. Horwitz J, Huang QL, Ding L, Bova MP. Lens alpha-crystallin: chaperone-like properties. *Methods Enzymol* 1998; 290:365-83. [PMID: 9534176]
 22. Takemoto LJ. Quantitation of asparagines -101 deamidation from alpha-crystallin during aging of the human lens. *Curr Eye Res* 1998; 17:247-50. [PMID: 9543632]
 23. Andley UP, Clark BA. Generation of oxidants in the near-UV photooxidation of human lens alpha-crystallin. *Invest Ophthalmol Vis Sci* 1989; 30:706-13. [PMID: 2703311]
 24. Jungblut PR, Otto A, Favor J, Löwe M, Müller EC, Kastner M, Sperling K, Klose J. Identification of mouse crystallins in 2D protein patterns by sequencing and mass-spectrometry. Application to cataract mutants. *FEBS Lett* 1998; 435:131-7. [PMID: 9762894]
 25. Ueda Y, Duncan MK, David LL. Lens proteomics: the accumulation of crystalline modifications in the mouse lens with age. *Invest Ophthalmol Vis Sci* 2002; 43:205-15. [PMID: 11773033]
 26. Mailankot M, Staniszevska MM, Butler H, Caprara MH, Howell S, Wang B, Doller C, Reneker LW, Nagaraj RH. Indoleamine 2,3-dioxygenase overexpression causes kynurenine-modification of proteins, fiber cell apoptosis and cataract formation in the mouse lens. *Lab Invest* 2009; 89:498-512. [PMID: 19308046]
 27. David LL, Azuma M, Shearer TR. Cataract and acceleration of calpain-induced beta-crystallin insolubilization occurring during normal maturation of rat lens. *Invest Ophthalmol Vis Sci* 1994; 35:785-93. [PMID: 8125740]
 28. Wang K, Gawinowicz MA, Spector A. The effect of stress on the pattern of phosphorylation of alphaA and alphaB crystallins in the rat lens. *Exp Eye Res* 2000; 71:385-93. [PMID: 10995559]
 29. Lampi KJ, Shih M, Ueda Y, Shearer TR, David LL. Lens proteomics: analysis of rat crystalline sequences and two-dimensional electrophoresis map. *Invest Ophthalmol Vis Sci* 2002; 43:216-24. [PMID: 11773034]
 30. Kador PF, Fukui HN, Fukushi S, Jernigan HM Jr, Kinoshita JH. Philly mouse: a new model of hereditary cataract. *Exp Eye Res* 1980; 30:59-68. [PMID: 7363969]
 31. Kuck JF. Late onset hereditary cataract of the emory mouse. A model for human senile cataract. *Exp Eye Res* 1990; 50:659-64. [PMID: 2197106]
 32. Okano T, Uga S, Ishikawa S, Shumiya S. Histopathological study of hereditary cataractous lenses in SCR strain rat. *Exp Eye Res* 1993; 57:567-76. [PMID: 8282043]
 33. Shumiya S. Establishment of the hereditary cataract rat strain (SCR) and genetic analysis. *Lab Anim Sci* 1995; 45:671-3. [PMID: 8746528]
 34. Mori M, Li G, Abe I, Nakayama J, Guo Z, Sawashita J, Ugawa T, Nishizono S, Serikawa T, Higuchi K, Shumiya S. Lanosterol synthase mutations cause cholesterol deficiency-associated cataracts in the Shumiya cataract rat. *J Clin Invest* 2006; 116:395-404. [PMID: 16440058]
 35. Kolosova NG, Lebedev PA, Aidagulova SV, Morozkova TS. OXYS rats as a model of senile cataract. *Bull Exp Biol Med* 2003; 136:415-9. [PMID: 14714098]
 36. Kolosova NG, Lebedev PA, Fursova AZh, Morozkova TS, Gusarevich OG. Prematurely aging OXYS rats as an animal model of senile cataract in human. *Adv Gerontol* 2003; 12:143-8. [PMID: 14743612]
 37. Marsili S, Salganik RI, Albright CD, Freel CD, Johnsen S, Peiffer RL, Costello M. Cataract formation in a strain of rats selected for high oxidative stress. *Exp Eye Res* 2004; 79:595-612. [PMID: 15500819]
 38. Solov'eva NA, Morozkova TS, Salganik RI. Development of a rat subline with symptoms of hereditary galactosemia and study of its biochemical characteristics. *Genetika* 1975; 11:63-71. [PMID: 1240812]
 39. Romyantseva, YuV; Fursova, AZh; Fedoseeva, LA.; Kolosova, NG. Changes in physicochemical parameters and α -crystallin expression in the lens during cataract development in OXYS rats. *Biochemistry (Mosc)* 2008; 73:1176-82. [PMID: 19120020]
 40. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72:248-54. [PMID: 942051]
 41. Ogden AT, Nunes I, Ko K, Wu S, Hines CS, Wang AF, Hegde RS, Lang RA. GRIFIN, a novel lens-specific protein related to the galectin family. *J Biol Chem* 1998; 273:28889-96. [PMID: 9786891]
 42. Jaworski C, Wistow G. LP2, a differentiation-associated lipid-binding protein expressed in bovine lens. *Biochem J* 1996; 320:49-54. [PMID: 8947466]
 43. Wen Y, Li GW, Chen P, Wong E, Bekhor I. Lens epithelial cell mRNA, II. Expression of a mRNA encoding a lipid-binding protein in rat lens epithelial cells. *Gene* 1995; 158:269-74. [PMID: 7607553]
 44. Ramaekers F, Dodemont H, Vorstenbosch P, Bloemendal H. Classification of rat lens crystallins and identification of proteins encoded by rat lens mRNA. *Eur J Biochem* 1982; 128:503-8. [PMID: 7151792]
 45. Robinson NE, Lampi KJ, Speir JP, Kruppa G, Easterling M, Robinson AB. Quantitative measurement of young human eye lens crystallins by direct injection Fourier transform ion cyclotron resonance mass spectrometry. *Mol Vis* 2006; 12:704-11. [PMID: 16807530]
 46. Srivastava OP, McEntire JE, Srivastava K. Identification of 9 kDa gamma-crystallin fragment in human lenses. *Exp Eye Res* 1992; 54:893-901. [PMID: 1521581]
 47. Andley UP. Crystallins in the eye: function and pathology. *Prog Retin Eye Res* 2007; 26:78-98. [PMID: 17166758]
 48. Shinohara T, Ikesugi K, Mulhern ML. Cataracts: role of the unfolded protein response. *Med Hypotheses* 2006; 66:365-70. [PMID: 16229964]

