

Heat shock protein 70 expression induced by diode laser irradiation on choroid-retinal endothelial cells in vitro

Shanshan Du, Qiong Zhang, Shisheng Zhang, Ling Wang, Jingcai Lian

Department of Ophthalmology, Ruijin Hospital, School of Medicine, Shanghai Jiaotong University, Shanghai, China

Purpose: To investigate the biologic effect of an 810 nm diode laser on the induction of heat shock protein 70 (Hsp70) in choroid-retinal endothelial cells in vitro.

Methods: Cultured rhesus macaque choroid-retinal endothelial cells (RF/6A) were irradiated using an 810 nm diode laser (spot size, 10 mm; duration, 60 s; power, 400–1,500 mW). Cell viability was assessed by annexin V- fluorescein isothiocyanate (FITC) and propidium iodide flow cytometric assay. Hsp70 expression was determined by western blot at 6, 12, 18, 24, and 48 h following laser exposure. Intracellular distribution of Hsp70 was examined by immunofluorescence staining.

Results: The laser-induced cell injury threshold was found to be at a power of 1,100 mW power (fluence, 84.08 J/cm²), above which there was significant cell death. Under this power, Hsp70 expression elevated obviously and was stronger at 600–1,000 mW power settings (fluences, 45.86–76.43 J/cm²). The expression of Hsp70 peaked at 12–18 h postirradiation, and returned to baseline by 48 h. Immunofluorescence staining indicated the induced Hsp70 expression in both the cytoplasm and the nucleus.

Conclusions: Subthreshold 810 nm diode laser exposure can induce Hsp70 hyperexpression from 12 to 18 h postirradiation in cultured choroid-retinal endothelial cells without obvious cell death. The results could be useful for investigating and designing more effective laser therapies.

Choroidal neovascularization (CNV) causes severe vision loss in several eye disorders such as age-related macular degeneration, ocular histoplasmosis, and high myopia [1]. Minimally invasive laser therapies have been commonly used, such as laser photocoagulation, photodynamic therapy (PDT), and transpupillary thermotherapy (TTT). Conventional laser photocoagulation is a well established treatment for some forms of CNV; nevertheless, it frequently damages the collateral neural retina, resulting in irreversible effects, including scotomata and the disruption of the retinal structure through scarring. PDT is an approach that specifically occludes subfoveal CNV without significant damage to adjacent tissues; however, a high proportion of secondary CNV and persistent choriocapillaris hypoperfusion have been found to occur with PDT and eventually lead to poor visual prognosis [2,3].

TTT, which is a long-exposure, large spot size, low-irradiance, near-infrared (diode 810 nm) laser protocol, slowly increases tissue temperature to approximately 4–10 °C above basal levels [4]. This procedure has been recently used with some success to slow down the progression of exudation and block subfoveal CNV in age-related macular degeneration [5-7]. However, clinical use of

TTT for CNV has been limited due to its lack of efficacy and occasional retinal damage [8,9]. It is difficult to devise an optimal laser parameter. Undertreatment would not close CNV, while overtreatment would lead to localized retinal destruction. Furthermore, the underlying mechanism of TTT remains unclear.

Heat shock proteins (Hsps) are a group of ubiquitous, highly conserved proteins in all organisms, and are classified according to their molecular mass. Elevated Hsps can be triggered by a variety of stressful stimuli and protect cells against stress. The 70 kDa Hsp (Hsp70) is particularly known to be induced by thermal, ischemic, and oxidative stress. Acting as a molecular chaperone, Hsp70 can assist in the refolding of denatured proteins and inhibit improper protein aggregation [10-12]. Moreover, the induction of Hsp70 may be responsible for the protective effect against apoptosis and inflammation [13-17]. A few animal experiments have found increased Hsp70 expression in the choroidoretinal layers [18-21] or optic nerve head [22] following TTT. From this perspective, TTT may be a subthreshold thermotherapy inducing Hsp70 hyperexpression, modulating apoptosis in choroidoretinal tissues [5,20,22,23]. Careful application of laser to induce Hsp70 hyperexpression may serve a protective role. Therefore, a better understanding of laser-induced Hsp70 kinetics is essential for the design of effective laser protocols. However, comprehensive Hsp70 expression kinetics data for choroidoretinal tissues or choroidoretinal related cells following TTT have not been reported.

Correspondence to: Ling Wang, Department of Ophthalmology, Ruijin Hospital, School of Medicine, Shanghai Jiaotong University, 197 2nd Ruijin Road, Shanghai, China 200025; Phone: 8613817230313; FAX: 86 21 64374104; email: lwang@rjeye.com



Figure 1. The setting for laser irradiation on RF/6A cells. Laser light was delivered through a 200 μm fiber-optic system, and the distance between the laser source and the cell monolayer was adjusted to obtain a laser spot that was 10 mm in diameter.

In this study, we focus on Hsp70 expression induced by 810 nm diode laser exposure in cultured choroid-retinal endothelial cells. We aim to elaborate on the conditions in which the protein is hyperexpressed, as well as the threshold at which cellular damage occurs.

METHODS

Cell culture: RF/6A rhesus macaque choroid-retinal endothelial cells were obtained from the Shanghai Institutes for Biologic Sciences (Shanghai, China) and were grown as monolayers in RPMI 1640 culture medium (Hyclone, Logan, UT) supplemented with 15% fetal bovine serum (Hyclone) plus 100 U/ml penicillin/streptomycin mixture (Sigma-Aldrich, Saint Louis, MO) in a 5% CO_2 humidified atmosphere at 37 $^\circ\text{C}$. Prior to laser irradiation, cells were seeded in 24-well culture plates (4×10^4 cells/well) to grow to reach about 80% confluence (about 3 days).

Laser irradiation: Irradiation was performed using an 810 nm diode laser (OcuLight SLx; Iridex Corporation, Mountain View, CA), delivered through a 200 μm fiber-optic system with different power outputs for 60 s in continuous wave mode. The distance between the laser source and the cell monolayer was adjusted to achieve a 10 mm diameter laser spot (Figure 1), adequate to cover about half a well of a 24-well plate. Before laser exposure, the medium was removed, the cells were washed once in sterile phosphate buffered saline (PBS; 137 mM NaCl, 5.4 mM KCl, 1.28 mM NaH_2PO_4 , 7 mM Na_2HPO_4 ; pH 7.4) and then transferred to no phenol red RPMI 1640 medium (Gibco, Grand Island, NY) without fetal bovine serum. The no phenol red medium was used to cover the cells during exposure to avoid absorption of laser energy by the culture

medium. Following irradiation, fresh culture medium was added and the cells were incubated for an additional period for measurements. The control group was treated in the same manner, except the laser irradiation. Another group without any treatment was defined as the normal group.

First, we observed the morphological change of RF/6A cells after laser exposure under an inverted microscope (Nikon TE2000-U; Nikon, Tokyo, Japan), and then identified the power required to cause significant cell death by annexin V-fluorescein isothiocyanate (annexin V-FITC) and propidium iodide (PI) flow cytometric assay (threshold application). Using a spot diameter of 10 mm and a power setting of 1,100 mW or greater, 60 s of laser irradiation produced significant cell death. Then, the induced Hsp70 expression was studied at various subthreshold power settings, using decrements of 100 mW (1,000, 900, 800, ... and 400 mW).

Detection of cell viability: Flow cytometry using Annexin V-FITC and PI double staining (BD Biosciences, San Diego, CA) was employed to detect cell viability. Following 6 h of incubation after exposure, cells were washed with PBS and resuspended in 60 μl binding buffer. annexin V-FITC (3.8 μl) and 3.8 μl PI were sequentially added to the cell solution. After incubation for 10 min in the dark, the stained cell solution was diluted with another 120 μl binding buffer and directly analyzed using the FACSCalibur system (BD Biosciences).

Western blot analysis: Based on the previous *in vivo* experimental data [18-20], we first detected Hsp70 expression with different subthreshold powers at 24 h postirradiation. Afterwards, the power resulting in stronger Hsp70 expression was used for observation of Hsp70 levels




Figure 2. Micrographs of RF/6A cells treated with different laser powers. **A:** Unirradiated RF/6A cells. **B:** RF/6A cells treated with 1,000 mW laser irradiation. **C:** RF/6A cells treated with 1,100 mW laser irradiation. Cells with apoptotic or necrotic morphology were present in the 1,100 mW treated samples. Data are from one of three separate experiments with similar results.

at 6, 12, 18, 24, and 48 h after irradiation. The cells were washed with ice-cold PBS and then lysed with radio immunoprecipitation assay (RIPA) buffer (Shenneng Bocai Biotechnology Co. Ltd, Shanghai, China) containing 1 mM phenylmethanesulfonyl fluoride (Shenneng Bocai Biotechnology Co. Ltd). The protein lysates were centrifuged at $12,000 \times g$ at 4°C for 30 min and the supernatant was collected for protein quantification using the bicinchoninic acid assay kit (Pierce, Rockford, IL). Equal amounts of protein (10–15 μg per lane) were separated by 10% sodium dodecyl sulfate PAGE and transferred onto a nitrocellulose membrane (Whatman Protran, Cambridge, UK). After blocking with 5% skimmed milk in 10 mmol/l Tris-buffered saline with 0.1% Tween-20 for 1 h, the blotted membrane was incubated with mouse monoclonal anti-Hsp70 antibody (1:1,000; Abcam, Cambridge, UK, no cross-reactivity to Hsc70) overnight at 4°C . After washing three times with 10 mmol/l Tris-buffered saline with 0.1% Tween-20, the membrane was incubated with IRDye 800CW goat anti-mouse secondary antibody (LI-COR, Lincoln, NE) for 1 h at room temperature. The signals were detected using an Odyssey Infrared Imaging System (LI-COR). Band intensity was analyzed by Quantity One software and compared with the internal standard glyceraldehyde-3-phosphate dehydrogenase (1:1,000; Abcam).

Immunofluorescence assay: Cells were prepared in triplicate on 0.1% gelatin-coated glass coverslips and treated with the laser, as described above. At 18 h after irradiation, adherent cells were fixed in 4% paraformaldehyde for 15 min, blocked with 10% goat serum containing 0.5% Triton X-100 in PBS for 30 min, and then incubated with mouse monoclonal anti-Hsp70 antibody (1:100; Abcam) overnight at 4°C . After three rinses with PBS, the samples were incubated with Cy3-conjugated goat anti-mouse secondary antibody (1:100, Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) for 1 h at room temperature in the dark. The slides were then washed in PBS and stained with 300 nM diaminodiphenylindole (Sigma-Aldrich, Saint

Louis, MO) for 10 min followed by three rinses with PBS. After mounting with fluorescent mounting media (Dako, Glostrup, Denmark), the slides were observed under a fluorescence microscope (Nikon 80i; Nikon).

Statistical analysis: All experiments were repeated at least three times. Statistical analysis was performed using SPSS 16.0 software. All error bars represent standard error of the mean (SEM). Differences between the laser-treated and control groups were evaluated by the Student paired *t* test. Statistical significance was defined as $p < 0.05$.

RESULTS

Inverted microscopy: The morphology of RF/6A cells treated with laser power lower than 1,100 mW was similar to that of the control (Figure 2A,B). However, a large number of cells with apoptotic or necrotic morphology (small, adherent or floating, rounded, or bubbling and shrunken) were present in samples treated with 1,100 mW laser power (Figure 2C).

Cell viability detection: We performed a flow cytometry experiment of double staining with Annexin V-FITC and PI to determine the cell viability and the laser injury threshold. The cells were distinguished into the following four groups (Figure 3A): those that were unlabeled (viable cells), those that were stained with annexin V-FITC only (early apoptotic), those that were stained with PI only (necrotic), and those that were stained with both annexin V and PI (late apoptotic/necrotic cells). Necrotic cells and the cells in early- and late-stage apoptosis were defined as dead.

In the control group, the percentage of viable cells was $95.54\% \pm 0.34\%$. After laser treatment with 900 mW, 1,000 mW, or 1,100 mW power, the percentages of viable cells were $94.54\% \pm 0.82\%$, $93.63\% \pm 0.50\%$, and $68.44\% \pm 2.66\%$, respectively (Figure 3B). Significant difference was observed between the 1,100 mW and the control, 900 mW, and 1,000 mW groups ($p = 0.008$, 0.009 , and 0.013 , respectively).

Western blot analysis: Figure 4 shows the western blot results of Hsp70 expression for both a variable laser power




Figure 3. Flow cytometry analysis for the levels of cell viability 6 h following laser irradiation using different powers based on annexin V-fluorescein isothiocyanate and propidium iodide double staining. **A:** Representative flow cytogram. The percentage of cells in the Q3 region denotes viable cells, Q4: early apoptotic, Q1: necrotic, and Q2: late apoptotic/necrotic cells. **B:** Graphical quantitation of the cell viability rates. The data represent mean \pm standard error of the mean (SEM) of three separate experiments (* $p < 0.05$ versus control cells, Student *t* test). The results revealed that the laser injury threshold was at a laser power of 1,100 mW, above which there was significant cell death.

with a fixed postirradiation period and a fixed laser power with a variable postirradiation period. The results demonstrated that Hsp70 was expressed at a very low level

in unirradiated cells. After laser exposure with a lower power (400–1,000 mW) followed by 18 h recovery, the protein expression elevated to different levels and was




Figure 4. Hsp70 expression of RF/6A cells after laser irradiation. **A:** RF/6A cells treated with various laser powers and followed by 18 h incubation. **B:** RF/6A cells treated with 700 mW laser power and followed by various incubation times. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as an internal control. Data are the mean±standard error of the mean (SEM) of three separate experiments (* $p < 0.05$ versus control cells, Student *t* test).

stronger at 600–1,000 mW power settings (fluences, 45.86–76.43 J/cm², Figure 4A).

We employed 700 mW (fluence, 53.50 J/cm²) laser power for observation of Hsp70 expression at different recovery times. Hsp70 expression at 1 and 3 h after laser exposure were similar to that of control (data not shown). The maximum expression of Hsp70 after laser exposure occurred at 12–18 h (about three times higher than the basal level), and gradually dropped to baseline at 48 h. Statistical analysis exhibited that Hsp70 expressions at 12, 18, and 24 h were all significantly higher than control ($p < 0.05$, Figure 4B).

Immunofluorescence assay: Immunofluorescence was performed to determine the expression and localization of induced Hsp70 in RF/6A cells. Cells were irradiated with 700 mW power, as already described, and incubated for 18 h. Unirradiated cells were used as control. As shown in Figure 5, Hsp70 was expressed at basal level in the nucleus and cytoplasm in unirradiated cells. After 700 mW laser exposure, a strong increase of Hsp70 was induced in both the cytoplasm and nucleus.

DISCUSSION

Laser irradiation at 810 nm, which was minimally absorbed by clear ocular media, was mostly delivered to retinal pigment epithelial and choroidal melanin. The absorption of the radiation could cause a limited temperature rise that may achieve occlusion of CNV [23,24]. It has recently been suggested that TTT could limit inflammation and play a major role in apoptosis and Hsp70 expression [23]. Hsp70 is necessary to make cells evolve toward survival instead of apoptosis or toward apoptosis instead of necrosis [25].

In effort to understand the laser-tissue interactions, some animal models have been investigated [18–21,26]. However, further biochemical and molecular analyses are not easily conducted through animal studies. Therefore, it has been difficult to devise an optimal therapeutic method based on a thorough understanding of the underlying mechanisms. In this study, we designed a laser-induced cell damage model to facilitate biologic research at the molecular level. To study the biologic effects on the cells, it is important to ensure that more cells are irradiated. Since the small spot diameters of commonly used lasers limit the number of cells irradiated, we selected a spot size of 10 mm




Figure 5. Expression and distribution of heat shock protein 70 in RF/6A cells after laser irradiation. **A-C**: Unirradiated RF/6A cells. **D-F**: RF/6A cells were treated with 700 mW laser irradiation and followed by 18 h incubation. Cells were fluorescently labeled with heat shock protein 70 (Hsp70; center, red channel) and the nuclear marker diamidinophenylindole (left, blue channel). The right is an overlay of the two channels. A strong induction of Hsp70 was observed in cells treated with a laser power of 700 mW. The induced Hsp70 was present in both the cytoplasm and nucleus. Data are from one of three separate experiments with similar results.

in diameter to cover about half a well, so as to receive the most prominent cell responses.

In animal models, the laser power that induced retinal whitening was thought to be the photocoagulation threshold [19-21]. In our *in vitro* experiment, the laser-induced cell injury threshold was determined by annexin V-FITC and PI flow cytometric assay. Significant cell death was found at a laser power of 1,100 mW (fluence, 84.08 J/cm²). Moreover, the results showed that 1,100 mW power induced about 31.56%±2.66% cell death. As the laser spot in our experiment only covered half a well, the actual percentage of 1,100 mW laser induced cell death could be more than 50%. In addition, the inverted microscopy showed a large number of apoptotic or necrotic cells in samples treated with 1,100 mW power. Therefore, we assumed that this laser power of 1,100 mW (fluence, 84.08 J/cm²) corresponded to the laser injury threshold.

After the cellular damage threshold was identified, a range of subthreshold powers from 400 to 1,000 mW was applied for detection of Hsp70 expression (with fluences ranging from 30.57 to 76.43 J/cm²). With these parameters, Hsp70 expression was highly induced, especially at 600–1,000 mW power settings (fluences, 45.86–76.43 J/cm²). As indicated by immunofluorescence staining, induced

expression of Hsp70 was present in both the cytoplasm and nucleus. Although the exposure conditions for Hsp70 induction in cultured cells in our study differed from those observed *in vivo*, their proportions of fluences for Hsp70 hyperexpression to fluences for the injury threshold are similar, about 60%–80% [19,20,26].

Hsp70 expression in our study was tracked for 48 h following laser exposure. The results showed that peak Hsp70 expression occurred at 12–18 h postirradiation, and reverted to baseline at 48 h, similar to the observations of Wang et al. [27], who used a matrix of heating and recovery times to determine the Hsp70 expression kinetics of bovine aortic endothelial cells in response to thermal stress. Kinetics data of Hsp70 expression following TTT *in vivo* have rarely been reported. In animal models, Hsp70 expression was mainly studied at 24 h after TTT.

The mechanisms of Hsp70 induction following 810 nm laser irradiation and the biologic action of Hsp70 merit further investigation. Since our RF/6A cells contained little melanin, they absorbed minimally with laser exposure, indicating that the induction of Hsp70 after laser exposure may be due to some nonthermal effects.

There were some limitations in our study. The laser spots were large in diameter and performed one by one,

which may have led to nonuniform energy distribution and a nonuniform cellular response. Raster scanning may produce homogeneous exposure of cells and yield uniform cellular response.

In summary, our study demonstrated that an 810 nm diode laser with subthreshold power (fluence lower than 84.08 J/cm²) applied for 60 s can stimulate Hsp70 production in cultured choroid-retinal endothelial cells without obvious cell damage. The Hsp70 expression was found to peak at 12 to 18 h following laser exposure, and returned to baseline at about 48 h. These results may facilitate designing more effective laser therapies.

ACKNOWLEDGMENTS

Supported by the National Natural Science Foundation of China (Grant No. 60,978,030) and the Shanghai Leading Academic Discipline Project (S30205). We thank Laboratory of Clinical Visual Sciences, Shanghai Institute of Health Sciences for technical help. None of the authors has any financial or proprietary interest in the materials or methods mentioned.

REFERENCES

- Grossniklaus HE, Green WR. Choroidal neovascularization. *Am J Ophthalmol* 2004; 137:496-503. [PMID: 15013874]
- Colucciello M. Choroidal neovascularization complicating photodynamic therapy for central serous retinopathy. *Retina* 2006; 26:239-42. [PMID: 16467692]
- Yaman A, Arikan G, Saatci AO, Cingil G. Choroidal neovascularization following photodynamic therapy in a patient with chronic central serous chorioretinopathy. *Bull Soc Belge Ophtalmol* 2007; 303:69-73. [PMID: 17894291]
- Ritchie KP, Keller BM, Syed KM, Lepock JR. Hyperthermia (heat shock)-induced protein denaturation in liver, muscle and lens tissue as determined by differential scanning calorimetry. *Int J Hyperthermia* 1994; 10:605-18. [PMID: 7806918]
- Newsom RS, McAlister JC, Saeed M, McHugh JD. Transpupillary thermotherapy (TTT) for the treatment of choroidal neovascularisation. *Br J Ophthalmol* 2001; 85:173-8. [PMID: 11159481]
- Ahuja RM, Benner JD, Schwartz JC, Butler JW, Steidl SM. Efficacy of transpupillary thermotherapy (TTT) in the treatment of occult subfoveal choroidal neovascularization in age-related macular degeneration. *Semin Ophthalmol* 2001; 16:81-5. [PMID: 15491008]
- Hogan AC, Kilmartin DJ. Low power vs standard power transpupillary thermotherapy in patients with age-related macular degeneration and subfoveal choroidal neovascularization ineligible for photodynamic therapy. *Eye (Lond)* 2006; 20:649-54. [PMID: 16082398]
- Mason JO 3rd, Colagross CC, Feist RM, White MF, Thomley ML, Vail RS, McGwin G Jr. Risk factors for severe vision loss immediately after transpupillary thermotherapy for occult subfoveal choroidal neovascularization. *Ophthalmic Surg Lasers Imaging* 2008; 39:460-5. [PMID: 19065975]
- Tsai TH, Yang CM, Yang CH, Ho TC, Huang JS, Chen MS. Transpupillary thermotherapy for the treatment of choroidal neovascularization in age-related macular degeneration in Taiwan. *Eye (Lond)* 2007; 21:721-6. [PMID: 16543926]
- Beckham JT, Wilmink GJ, Mackanos MA, Takahashi K, Contag CH, Takahashi T, Jansen ED. Role of HSP70 in cellular thermotolerance. *Lasers Surg Med* 2008; 40:704-15. [PMID: 19065555]
- Rylander MN, Feng Y, Bass J, Diller KR. Thermally induced injury and heatshock protein expression in cells and tissues. *Ann N Y Acad Sci* 2005; 1066:222-42. [PMID: 16533928]
- Lanneau D, de Thonel A, Maurel S, Didelot C, Garrido C. Apoptosis versus cell differentiation: role of heat shock proteins HSP90, HSP70 and HSP27. *Prion* 2007; 1:53-60. [PMID: 19164900]
- Didelot C, Schmitt E, Brunet M, Maingret L, Parcellier A, Garrido C. Heat shock proteins: endogenous modulators of apoptotic cell death. *Handb Exp Pharmacol* 2006; 172:171-98. [PMID: 16610360]
- Söti C, Nagy E, Giricz Z, Vigh L, Csermely P, Ferdinandy P. Heat shock proteins as emerging therapeutic targets. *Br J Pharmacol* 2005; 146:769-80. [PMID: 16170327]
- Franklin TB, Krueger-Naug AM, Clarke DB, Arrigo AP, Currie RW. The role of heat shock proteins Hsp70 and Hsp27 in cellular protection of the central nervous system. *Int J Hyperthermia* 2005; 21:379-92. [PMID: 16048836]
- Awasthi N, Wagner BJ. Upregulation of heat shock protein expression by proteasome inhibition: an antiapoptotic mechanism in the lens. *Invest Ophthalmol Vis Sci* 2005; 46:2082-91. [PMID: 15914627]
- Yenari MA, Liu J, Zheng Z, Vexler ZS, Lee JE, Giffard RG. Antiapoptotic and anti-inflammatory mechanisms of heat-shock protein protection. *Ann N Y Acad Sci* 2005; 1053:74-83. [PMID: 16179510]
- Desmettre T, Maurage CA, Mordon S. Transpupillary thermotherapy (TTT) with short duration laser exposures induce heat shock protein (HSP) hyperexpression on choroidoretinal layers. *Lasers Surg Med* 2003; 33:102-7. [PMID: 12913881]
- She H, Li X, Yu W. Subthreshold transpupillary thermotherapy of the retina and experimental choroidal neovascularization in a rat model. *Graefes Arch Clin Exp Ophthalmol* 2006; 244:1143-51. [PMID: 16429304]
- Desmettre T, Mauragr CA, Mordon S. Heat shock protein hyperexpression on chorioretinal layers after transpupillary thermotherapy. *Invest Ophthalmol Vis Sci* 2001; 42:2976-80. [PMID: 11687545]
- Morimura Y, Okada AA, Hayashi A, Fujioka S, Hashida N, Kawahara S, Hida T. Histological effect and protein expression in subthreshold transpupillary thermotherapy in rabbit eyes. *Arch Ophthalmol* 2004; 122:1510-5. [PMID: 15477463]
- Kim JM, Park KH, Kim YJ, Park HJ, Kim DM. Thermal injury induces heat shock protein in the optic nerve head in vivo. *Invest Ophthalmol Vis Sci* 2006; 47:4888-94. [PMID: 17065503]
- Mainster MA, Reichel E. Transpupillary thermotherapy for age-related macular degeneration, long-pulse photocoagulation, apoptosis, and heat shock proteins. *Ophthalmic Surg Lasers* 2000; 31:359-73. [PMID: 11011704]

24. Reichel E, Berrocal AM, Ip M, Kroll AJ, Desai V, Duker JS, Puliafito CA. Transpupillary thermotherapy of occult subfoveal choroidal neovascularization in patients with age-related macular degeneration. *Ophthalmology* 1999; 106:1908-14. [PMID: 10519584]
25. Samali A, Orrenius S. Heat shock proteins: regulators of stress response and apoptosis. *Cell Stress Chaperones* 1998; 3:228-36. [PMID: 9880235]
26. Sramek C, Mackanos M, Spitler R, Leung LS, Nomoto H, Contag CH, Palanker D. Non-damaging retinal phototherapy: dynamic range of heat shock protein expression. *Invest Ophthalmol Vis Sci* 2011; 52:1780-7. [PMID: 21087969]
27. Wang S, Diller KR, Aggarwal SJ. Kinetics Study of Endogenous Heat Shock Protein 70 Expression. *J Biomech Eng* 2003; 125:794-7. [PMID: 14986403]

Articles are provided courtesy of Emory University and the Zhongshan Ophthalmic Center, Sun Yat-sen University, P.R. China.

The print version of this article was created on 20 September 2012. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.