

Supplementary Figure 1

TGF β -induced IOP elevations are mediated by RhoA in the early but not the late fibrotic phase of open angle glaucoma

Lisa J Hill, PhD^{a,1,*}, Ben Mead, PhD^{b,1}, Chloe Thomas, BMedSc^a, Simon Foale BMedSc^a, Elena Feinstein, PhD^c, Martin Berry, DPhil^a, Richard J Blanch, PhD^a, Zubair Ahmed, PhD^{a,2} and Ann Logan, PhD^{a,2}

^aNeuroscience and Ophthalmology Research Group, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, B15 2TT, United Kingdom

^bSection of Retinal Ganglion Cell Biology, Laboratory of Retinal Cell and Molecular Biology, National Eye Institute, National Institutes of Health, Bethesda, Maryland, 20892, USA

^cResearch Division, Quark Pharmaceuticals, Ness Ziona, Israel

¹Contributed equally and are joint first authors

²Joint senior authors

* **Corresponding author:** Lisa Hill, Neuroscience and Ophthalmology Research Group, Institute of Inflammation and Ageing, University of Birmingham, B15 2TT, United Kingdom. Tel: +44(0) 121 414 8859. Email address: L.J.Hill@bham.ac.uk

