



The Cloning of GRK7, a Candidate Cone Opsin Kinase, from Cone- and Rod-Dominant Mammalian Retinas

Ellen R. Weiss,¹Dayanidhi Raman,¹Satoko Shirakawa,¹ Melissa H. Ducceschi,¹ Paul T. Bertram,¹Fulton Wong,²Timothy W. Kraft,³Shoji Osawa¹

¹Department of Cell Biology and Anatomy, The University of North Carolina at Chapel Hill, Chapel Hill, NC; ²Department of Ophthalmology, Duke University Eye Center, Durham, NC; ³Department of Ophthalmology, University of Alabama, Birmingham, AL

Purpose: Desensitization in the rod cell of the mammalian retina is initiated when light-activated rhodopsin is phosphorylated by the G protein-coupled receptor kinase (GRK). GRK1, often referred to as rhodopsin kinase. A distinct kinase that specifically phosphorylates cone opsins in a similar manner has not been identified in mammals. To determine the existence of a cone opsin kinase, RNA from the retinas of cone- and rod-dominant mammals was analyzed by PCR.

Methods: RNA prepared from the retinas of two cone-dominant mammals, the thirteen-lined ground squirrel and the eastern chipmunk, and a rod-dominant mammal, the pig, was used to clone a new GRK family member by RT-PCR. The tissue distribution and localization of the kinase in retina were determined by Northern blot hybridization and in situ hybridization. The protein encoded by this cDNA was expressed in human embryonic kidney-293 (HEK-293) cells and compared with bovine GRK1 for its ability to phosphorylate bovine rhodopsin and to undergo autophosphorylation.

Results: The cDNA cloned from ground squirrel contains an open reading frame encoding a 548 amino-acid protein. Sequence analysis indicates that this protein is orthologous to GRK7 recently cloned from *O. latipes*, the medaka fish. Partial cDNA fragments of GRK7 were also cloned from RNA prepared from eastern chipmunk and pig retinas. In situ hybridization demonstrated widespread labeling in the photoreceptor layer of the ground squirrel retina, consistent with expression in cones. Recombinant ground squirrel GRK7 phosphorylates bovine rhodopsin in a light-dependent manner and can be autophosphorylated, similar to bovine GRK1.

Conclusions: These results indicate that cone- and rod-dominant mammals both express GRK7. The presence of this kinase in cones in the ground squirrel and its ability to phosphorylate rhodopsin suggests that it could function in cone cells as a cone opsin kinase.

In the rod cell of the mammalian retina, GRK1, a G protein-coupled receptor kinase (GRK) also referred to as rhodopsin kinase, phosphorylates light-activated rhodopsin and promotes the binding of arrestin to terminate visual signaling by transducin (G_t), the rod cell G protein [1]. Analogous events involved in cone visual signaling are less well-characterized, due to the fact that mammalian retinas generally have many more rods than cones. The cloning of a number of cone-specific proteins with significant homology to their rod cell counterparts [2-12] suggests that the signaling pathway is similar. However, dramatic physiological differences are observed between rods and cones in their responses to light. For example, cones are typically orders of magnitude less sensitive to similar light intensities, their response time is more rapid, and signal termination is faster than in rods [13]. Several mechanisms may account for the difference in the rate of signal termination. For example, the decay of the active forms (meta II) of the cone opsins is faster than the decay of rhodopsin [14,15]. It also seems likely that the lifetime of activated transducin is different in rods and cones, given the new evidence for regulation of GTP hydrolysis of the G protein α subunits by members of the Regulators of G Protein Signal-

ing (RGS) family. RGS9, which has been shown in vitro to accelerate the rate of GTP hydrolysis for α_{11} , the rod cell transducin α subunit, is actually present at higher levels in cones [16,17]. In addition, differences in the kinetics of opsin phosphorylation and arrestin binding could play a role in the different rates of signal termination observed for rods and cones.

Six GRKs have been cloned from mammals and biochemically characterized [18,19]. Recently, a novel member of the GRK family, GRK7, was cloned from the medaka fish (*Oryzias latipes*) and was localized specifically to cones, indicating that the cones of lower vertebrates have a distinct GRK [20]. Whether a unique GRK phosphorylates the cone opsins and plays a role in the termination of visual signaling in mammals is unknown. GRK1 has been localized to both rod and cone outer segments in mammals by immunohistochemistry [21], raising the possibility that GRK1 acts to desensitize both rod and cone opsins. However, in patients with Oguchi disease, which is caused by inactivating mutations in GRK1 [22,23], cone responses to light are relatively normal [24], suggesting that GRK1 does not play a major role in cone visual signaling.

The present study describes the cloning of a new mammalian member of the GRK family from the 13-lined ground squirrel and the eastern chipmunk, two cone-dominant mammals, as well as from the pig, a rod-dominant mammal. Sequence analysis suggests that this kinase is mammalian GRK7. The mRNA for this GRK is found exclusively in the retina and only in the photoreceptor cell layer of the ground squirrel

Correspondence to: Shoji Osawa, Ph.D., Department of Cell Biology and Anatomy, The University of North Carolina at Chapel Hill, CB# 7090, 108 Taylor Hall, Chapel Hill, NC, 27599-7090; Phone: (919) 966-7683; FAX: (919) 966-1856; email: shoosawa@med.unc.edu

retina. Like medaka fish GRK7, the 13-lined ground squirrel GRK7 contains a consensus sequence for geranylgeranylation of the C terminus. Functional studies demonstrate that this kinase phosphorylates bovine rhodopsin in a light-dependent manner and can be autophosphorylated, similar to GRK1. In addition to GRK7, we successfully cloned a partial cDNA fragment of GRK1 from both ground squirrel and pig retina RNA. These results support the existence of a new GRK family member (GRK7) in both cone- and rod-dominant mammalian retinas that is distinct from GRK1 and has the potential to regulate desensitization in cones.

METHODS

Preparation of RNA: Thirteen-lined ground squirrels were purchased from TLS Research (Bartlett, IL). Eastern chipmunks were obtained by trapping the animals in Jefferson County, Alabama. The animals were euthanized by carbon dioxide asphyxiation. The eyes were enucleated and the retinas frozen in liquid nitrogen. Animal care guidelines in accordance with those published by the Institute for Laboratory Animal research were followed. Total RNA from ground squirrel and eastern chipmunk retinas was prepared using the TRIzol Reagent (Life Technologies, Gaithersburg, MD) according to the manufacturer's protocols. Poly(A)⁺ RNA was prepared using the polyAtract Kit (Promega, Madison, WI) following procedures described by the manufacturer. Rat poly(A)⁺ RNA was purchased from Clontech (Palo Alto, CA).

Cloning of GRK7: Two degenerate primers used previously to clone members of the GRK family [25-29], were modified to recognize all GRKs and used to amplify sequence from rat, 13-lined ground squirrel, and eastern chipmunk retina RNA. Poly(A)⁺ RNA or total RNA from rat, eastern chipmunk, and thirteen-lined ground squirrel were reverse-transcribed with MMLV reverse transcriptase using either a random hexamer or oligo dT primer (Advantage RT-for-PCR Kit, Clontech). Polymerase chain reaction (PCR) was performed with Pfu polymerase (Stratagene, La Jolla, CA) using two degenerate oligonucleotide primers, GT(ACGT)TA(CT)(AC)G(ACGT)GA(CT)(CT)T(ACGT)AA(AF)CC and A(ACG)(CT)TC(ACGT)GG(ACGT)GCCAT(AG)(AT)A(ACGT)CC, that correspond to amino acids 311-317 (I/VYRDLKP) and 354-360 (GF/YMAPEL/V) of bovine GRK1 within a conserved region of the catalytic domain. A 2-min incubation at 95 °C was followed by 30-35 amplification cycles, each containing 1 min at 95 °C, 1 min at 50 °C and 2 min at 72 °C. An approximately 150-bp fragment isolated on an agarose gel was ligated to pCR-script SK (Stratagene) and sequenced using the dideoxy sequencing method with the enzyme Sequenase (Amersham, Arlington Heights, IL). Alternatively, clones were sequenced at the UNC-CH Automated DNA Sequencing facility on a Model 377 DNA Sequencer (Perkin Elmer, Applied Biosystems Division, Foster City, CA) using the ABI PRISM™ Dye Terminator cycle Sequencing Ready Reaction Kit with AmpliTaq DNA Polymerase, FS (Perkin Elmer, Applied Biosystems Division).

The sequence of GRK7 from 13-lined ground squirrel RNA was extended using a Rapid Amplification of cDNA Ends (RACE) strategy [30]. For 3' extension, the 3' RACE System

kit (Life Technologies) was used according to the manufacturer's protocols. The GRK7 cDNA was amplified by PCR using two primers, AUAP (Abridged Universal Amplification Primer) and the primer GAGAACGTGCTCCTGGATGACCTCGG, which corresponds to a site (nucleotides 1040-1065) within the 146-bp sequence of GRK7 originally cloned by PCR. A 2.1-kb fragment generated using this method was ligated to pCR-script SK (Stratagene) and sequenced as described above.

For 5' extension of GRK7 from 13-lined ground squirrel RNA, the Marathon cDNA Amplification Kit (Clontech) was used. Approximately 2 µg of total RNA was reverse-transcribed by MMLV reverse transcriptase using the primer GGAATAGCTCGCCTTGTCCATCAGGAT, which corresponds to a site (nucleotides 1163-1189) downstream from the original 146-bp sequence of GRK7. Second strand synthesis was followed by ligation of Marathon cDNA adapters. Amplification of the cDNA of the GRK7 was performed by PCR using the gene-specific primer (described above) and the Adapter-Primer 2. A 1.2-kb fragment was purified and ligated to pCR2.1 (Invitrogen, Carlsbad, CA) and sequenced.

To verify the accuracy of the cDNA sequence, a minimum of 3 clones from different PCR reactions were analyzed for each region. Searches against peptide and nucleotide sequence databanks were performed at the GSC using the BLAST network services. Other sequence analyses were performed using programs from the Wisconsin Package Version 9.0 (Genetics Computer Group, Madison, WI) as described in the legends to Figs. 1 and 2. The nucleotide sequence for 13-lined ground squirrel GRK7 has been deposited in the GenBank database under the GenBank Accession Number [AF063016](#). The accession numbers of sequences found to be related to GRK7 are: [AB009568](#), *O. latipes* GRK7; [AB009569](#), *O. latipes* GRK1; [P28327](#), bovine GRK1; [Q15835](#), human GRK1; [P25098](#), human GRK2; [P35626](#), human GRK3; [P32298](#), human GRK4; [P34947](#), human GRK5; [P43250](#), human GRK6.

A partial fragment of the pig GRK7 cDNA was cloned by PCR using two degenerate oligonucleotide primers, GA(CT)TGGT(CT)GC(ACGT)ATGGG(ACGT)TG and TC(AGT)AT(CT)TC(AG)TC(ACGT)AC(AG)TC(CT)TT, corresponding to amino acids 368-374 and 476-482 of the ground squirrel sequence, respectively. A partial fragment of GRK1 was isolated from ground squirrel and pig using two degenerate primers, GA(AG)AA(AG)GT(ACGT)GA(AG)AA(CT)AA(AG)GA and TC(CT)TG(AG)AA(AG)AA(CT)TC(ACGT)GT(AG)TC, corresponding to amino acids 394-400 and 499-505 of bovine GRK1, respectively.

Northern blot hybridization analysis: Electrophoresis was performed following procedures described by Qiagen (Chatsworth, CA) in the RNeasy Mini Handbook. Approximately 8 µg of total RNA was electrophoresed on a formaldehyde-agarose gel and electrophoretically transferred to Immobilon-S nylon membrane (Boehringer Mannheim Indianapolis, IN). The RNA was immobilized by UV crosslinking. The membrane was prehybridized in 6X SSC, 5X Denhardt's solution, 0.5% SDS, 50% formamide and 100 µg/ml salmon sperm DNA at 42 °C for 2 h. A probe corresponding to the initial 146-bp fragment of GRK7 labeled with

A to remove unincorporated ATP, the protein was released from the resin by incubation with SDS-Laemmli buffer [32] and electrophoresed on 10% SDS-polyacrylamide gels. Phosphorylation of GRK7 was visualized by phosphorimager analysis of the dried gels.

RESULTS

Based on the hypothesis that a cone opsin kinase would be a member of the GRK family, a strategy was developed to clone new GRKs from the retinas of 2 cone-dominant mammals,

GRK7 (GS)	M.DMGGLDNL IANTAYLQAR KT.DSDSREL QRRRSLA.. .LPGPQGCAE	45
GRK7 (O1)	-C-----V-----K-Q ..GG-DK-M KK----S.. --K-EQ-VA	44
GRK1 (Bv)	..-F-S-ETV V--S-FIA-- GSF-AS-GPA S-D-KY--RL K--PLSK-EA	49
GRK7 (GS)	LRQSLSPHFH SLCEQQPIGR LRFDRFLATV PKYSQAVAFI EDVQNWELAE	95
GRK7 (O1)	--E-IEKD-T L--R---K -----NT -EFKL-AEF- DELYD-D---	94
GRK1 (Bv)	--E--DLG-E GM-LE---K -----QQ--R-H EQHGP-LQLW K-IEDYDT-D	99
GRK7 (GS)	EGPAKTSTLQ QLAATCARDP GP.QSFLSQD LATKCRRAST DEERKTLVEQ	144
GRK7 (O1)	GAAKDKARQN IINKY-KP-S KTFPLT---GE P-E--KSVTD ATFPEVMKNK	144
GRK1 (Bv)	DALRFPQKAQ LR--YLEPQA QLFCS---DAE TVARA--GAG -GLFQP-LR.	148
GRK7 (GS)	AKAETMSFLQ EQPFQDFLAS PFYDRFLQWK LFEMQPVS DK YFTEFRVLGK	194
GRK7 (O1)	VQDGVRE-K GK--TEYQG- QYF-K----- EY-K--I--- --Y--T---	194
GRK1 (Bv)	..AVLAH-G QA---E--D- LYFL----- WL-A--MGED W-LD--V--R	195
GRK7 (GS)	GGFGEVCAVQ VRNTGKMYAC KKLDKKRLKK KGGEKMALLE KEILEKVNSP	244
GRK7 (O1)	-----K-----Q-----C-----Q-----L	244
GRK1 (Bv)	-----F-C- MKA--Q-----N-----RK-YQG-MV- -K--A--H-R	245
GRK7 (GS)	FIVSLAYAFE SKTHLCLVMS LMNGDGLKFH IYVNG..... .TRGLAMSRV	288
GRK7 (O1)	-L-N-----YD T-----T-----Y-----I-YDGKG VDK-IE-K-I	294
GRK1 (Bv)	-----T--D-----T I-----IRY- ----DEDNP.FOEP-A	291
GRK7 (GS)	IFYTAQMTCS VHLHLGLGIV YRDLKPNVNL LDDLGNCRSL DLGLAVEVQD	338
GRK7 (O1)	-H-----I-T- I---DMD-I ---M-----SQ-Q-----I-IAP	344
GRK1 (Bv)	-----IVS- LE---QRN-I -----D--V-I- -----LKA	341
GRK7 (GS)	DKPITQ.RAG TNGYMAPEIL MDKASYSYPV DWFAMGCSIY EMVAGRTPFK	387
GRK7 (O1)	G-TV--.M- -GA-----S-TP-RTS- --W-L-----Y-----	392
GRK1 (Bv)	GQTK-KGY-- -P-F-----L- LGE.E-DFS- -Y--L-VTL- --I-A-G-R	390
GRK7 (GS)	..DFKEKVS KEDLKERTMK DEVAPHENF TEETKDCICRL FLAKKPEQRL	434
GRK7 (O1)	GPESK---E --EVQR-ILN E-PKWE-KC DAP--VIQQ -K--IDE--	442
GRK1 (Bv)	ARG.....E NKE--Q-VLE QA-TYPDK.- SPAS--F-EA L-Q-D--K--	436
GRK7 (GS)	GSRE.KADDP RKHPFFQTVN FPRLEAGLVE PPFVDPSPV YAKDVDEIDD	483
GRK7 (O1)	-M-N-NME-- --EW-KSI- -----D --W--K-N-- --K--TGD-AE	491
GRK1 (Bv)	-F-DGSC-GL -T--L-RDIS WRQ---MLT -----SRT- ----NIQDVGA	486
GRK7 (GS)	FSEVRGVEFD DKDKQFFQRF STGAVPVAWQ EEIIEIETGLFE ELN.....	526
GRK7 (O1)	---IK-I--- A--DK--KE- -----IQ-- Q-M-----L-D -----	534
GRK1 (Bv)	--T-K--A-E KA-TE--E- AS-TC-IP-- --M-----V-G D--VWRPDGQ	536
GRK7 (GS)	..DPNRPSGD GKG.DSSKSG VCLLL 548	
GRK7 (O1)	..---KE-A -G-D-EK--- T-A-- 557	
GRK1 (Bv)	MP-DMKGVSG QEAAPE---- M-V-S 561	

Figure 2. Comparison of the amino-acid sequence of ground squirrel GRK7 to GRK7 from *Oryzias latipes* and bovine GRK1. The hyphens (-) represent sequence in GRK7 from *Oryzias latipes* (O1) and bovine (Bv) GRK1 that is identical with the sequence of ground squirrel (GS) GRK7. Gaps in the sequence alignment are represented by periods (.). The blue lettered region represents the predicted catalytic domain (amino acids 185-464 of the ground squirrel sequence). The asterisks (*) mark serine and threonine residues that serve as known or potential autophosphorylation sites. The sequence with the double underline represents the CAAX motif, which codes for isoprenylation at the cysteine residue. Sequence comparisons were performed using the computer programs PILEUP and PRETTY from the Wisconsin Package Version 9.0. The nucleotide sequence for 13-lined ground squirrel GRK7 has been deposited in the GenBank database under the GenBank Accession Number AF063016.

the eastern chipmunk [37] and the 13-lined ground squirrel [38-39]. PCR primers corresponding to amino acids 311-317 and 354-360 of bovine GRK1 were designed to recognize the known GRK family members and used to screen retinal RNA from these animals. The cloned PCR products were classified according to their sequence and compared with the products from a PCR analysis of RNA from the rat, a rod-dominant animal [40]. As expected, the majority (85%) of GRKs cloned from rat retina RNA were GRK1 (data not shown). GRK1 was not detected in the eastern chipmunk or the 13-lined ground squirrel during this initial screening. However, a 146-bp fragment corresponding to a novel member of the GRK family was cloned from both the ground squirrel and eastern chipmunk. The sequences from these two animals were 91% identical at the amino acid level, excluding the primer hybridization sites, suggesting that they represent orthologous proteins (species homologues) (Figure 1A).

A RACE strategy was used to obtain the 5' and 3' regions of the clone from ground squirrel; 2.1 kb and 1.2 kb fragments, respectively, were obtained that overlapped each other. The full-length cDNA is 3095 bases and contains a single open reading frame from nucleotides 98 to 1744 encoding a polypeptide of 548 amino acids. The calculated molecular mass of the protein is 61,993 Da. The C terminus contains a CAAX motif, coding for posttranslational attachment of a geranylgeranyl group to Cys-545. Alignment of this sequence with the recently cloned GRK7 from *O. latipes* (Figure 2) demonstrated an overall identity between these two proteins of 59%. In contrast, the ground squirrel GRK was found to be only 48% identical with bovine GRK1. The highly conserved catalytic domains of the GRK family members, consisting of approximately 280 amino acids, were also compared (Table 1). The catalytic domain of the ground squirrel kinase shows greatest identity with *O. latipes* GRK7 (67%), is more distantly related to GRK1, 4, 5 and 6 (approximately 55-57% identity), and shows the least sequence identity with GRK2 and GRK3 (approximately 40% identity). These results suggest that the ground squirrel GRK is a new mammalian GRK and orthologous to GRK7 cloned from *O. latipes*. Despite the similarity between ground squirrel GRK7 and GRK1, 4, 5 and 6, analysis of the evolutionary distance between these proteins

TABLE 1. HOMOLOGY BETWEEN THE CATALYTIC DOMAINS OF GRK7 AND OTHER MEMBERS OF THE GRK FAMILY.

	GRK7 (GS)	
	Identity (%)	Similarity (%)
GRK7 (GS)	100.0	100.0
GRK7 (<i>O. latipes</i>)	67.4	77.1
GRK1 (bovine)	55.4	66.2
GRK1 (human)	56.5	65.8
GRK2 (human)	40.4	52.4
GRK3 (human)	39.6	50.9
GRK4 (human)	56.8	67.3
GRK5 (human)	56.5	68.3
GRK6 (human)	56.1	67.3

Protein sequence data was analyzed using the computer program GAP from the Wisconsin Package version 9.0

indicates that this GRK7 is most closely related to GRK1 (not shown), similar to the observations of Hisatomi et al. for *O. latipes* GRK7 [20].

RT-PCR was used to amplify GRK7 sequence from RNA isolated from the pig retina. The pig retina is rod-dominant, composed of 89% rods and 11% cones [41,42]. Two degenerate oligonucleotide primers, based on the sequence of ground squirrel GRK7, were used to clone a 303-bp fragment from pig RNA corresponding to amino acids 375-475 of the ground squirrel protein (Figure 1B). The identity of this region between ground squirrel and pig GRK7 is 84%. In addition, partial fragments of GRK1 were also cloned from both ground squirrel and pig (Figure 1C). This region of ground squirrel GRK1 is 79% and 83% identical to pig and bovine GRK1, respectively. An overlapping region of the GRK7 and GRK1 partial cDNAs, consisting of 78 amino acids, demonstrates only 43% identity to each other in ground squirrel and 47% identity in the pig. Therefore, GRK7 and GRK1 are distinct kinases present in both rod- and cone-dominant mammals.

The distribution of GRK7 in 8 ground squirrel tissue RNA preparations was determined using Northern blot hybridization (Figure 3). A probe corresponding to the original 146-bp sequence of GRK7 hybridized only to retinal RNA. The molecular size of the major band is approximately 3 kb, which corresponds closely to the length of the complete cDNA sequence (3095 bp) of GRK7. Several minor bands are visible above and below the major band. Some of these may be processed forms, or splice variants, of GRK7, such as those reported for GRK1 and GRK4 [21,43,44]. These results suggest that ground squirrel GRK7 is highly tissue-specific in its expression and is found predominantly in the retina. The same probe was used to localize GRK7 messenger RNA by in situ hybridization in histological preparations of ground squirrel




Figure 3. Northern blot analysis of GRK7. A ^{32}P -labeled probe of GRK7 was hybridized to a blot containing ground squirrel tissue RNA as described in the "Methods". Lane 1: brain, Lane 2: heart, Lane 3: kidney, Lane 4: liver, Lane 5: lung, Lane 6: retina, Lane 7: skeletal muscle, Lane 8: spleen. Numbers at left represent molecular size markers for RNA (Life Technologies).

retina (Figure 4). The antisense-strand RNA probe hybridized to the photoreceptor cell layer, which is composed of approximately 94% cones and 6% rods [38]. In contrast, the sense-strand RNA probe showed no specific hybridization. These results indicate that GRK7 is distributed exclusively in photoreceptor cells of the 13-lined ground squirrel retina, consistent with a function in cone visual signaling.

To determine whether the cDNA for ground squirrel GRK7 encodes a protein of the correct molecular size, the cDNA was expressed by in vitro translation. The [^{35}S]methionine-labeled, in vitro-translated product of the GRK7 cDNA migrates at approximately 62 kDa, consistent with the molecular size of the protein predicted from the length of the open reading frame (data not shown). Ground squirrel GRK7 and bovine GRK1 were also transiently expressed in HEK-293 cells to compare their ability to phosphorylate rhodopsin. Cytosolic extracts prepared from these cells were analyzed for their ability to phosphorylate bovine rhodopsin in ROS membranes in the light and in the dark (Figure 5). GRK7 and GRK1 were similarly effective at phosphorylating rhodopsin in a light-dependent manner. Extracts prepared from nontransfected cells showed no detectable phosphorylation of rhodopsin. These results indicate that GRK7 can phosphorylate rhodopsin in a light-dependent manner similar to GRK1, consistent with its classification as a GRK family member and with a potential role as a cone opsin kinase.

Autophosphorylation sites have been identified for GRK1, 5, and 6 downstream from the catalytic domain [28,45-48]. Autophosphorylation of GRK1 may regulate the binding of ATP to the catalytic domain and the selectivity for different substrate sites on rhodopsin [46]. Ground squirrel GRK7 has a threonine at position 21, equivalent to a minor autophosphorylation site at Ser-21 in bovine GRK1 (Figure 2). A serine is also present at position 485, which is equivalent to one of two major autophosphorylation sites at Ser-488 in bovine GRK1 [46], suggesting that GRK7 may also be autophosphorylated. A histidine-tagged GRK7 construct expressed in HEK-293 cells was examined for its ability to un-




Figure 4. Localization of GRK7 mRNA in ground squirrel retinas using in situ hybridization. ^{35}S -labeled antisense and sense RNA probes were produced from the 146-bp fragment corresponding to nucleotide sequence 1019-1164 by in vitro transcription and hybridized to cryostat sections of ground squirrel retina as described in the "Methods". (A) Dark field image of ground squirrel retina hybridized with an antisense probe for GRK7. Retinal pigment epithelium (RPE), photoreceptor cell layer (PCL), inner segment layer (INL), ganglion cell layer (GCL). (B) Dark field image of ground squirrel retina hybridized with a sense probe for GRK7. Abbreviations are as described in (A). Since the retinal pigment epithelium is black, it appears white in dark field images.

dergo autophosphorylation. Extracts from transfected and nontransfected cells were partially purified on Ni-NTA resin and incubated with [γ^{32} P]ATP (Figure 6). The results using extracts from transfected cells show the presence of a radioactive protein bound to the resin with a molecular size similar to that of GRK7 identified by *in vitro* translation. In contrast, this band was not detected in 32 P-labeled extracts from nontransfected cells or in the flowthrough from either transfected or nontransfected cell extracts. These results suggest that GRK7, like GRK1, GRK5 and GRK6, is capable of autophosphorylation.

DISCUSSION

The present report describes the cloning of a new member of the mammalian GRK family from the retinas of the 13-lined ground squirrel, the eastern chipmunk and the pig. Comparison of this sequence with known GRK family members and the sequence of the newly cloned GRK7 from *O. latipes*, the medaka fish, suggests that these proteins are mammalian GRK7. Although its sequence is closely related to GRK1, partial fragments of a different cDNA cloned from the 13-lined ground squirrel and pig show greater homology to GRK1. Therefore cone- and rod-dominant mammals both express GRK1 and GRK7. Northern analysis and *in situ* hybridization demonstrate that GRK7 is expressed exclusively in the photoreceptor cell layer of the ground squirrel. Approximately 94% of the photoreceptors of the ground squirrel retina are cones [38], making GRK7 an excellent candidate for a cone opsin kinase. Furthermore, *in vitro* phosphorylation studies show that rhodopsin can serve as a substrate for this kinase, strengthening the likelihood that it can phosphorylate the homologous cone opsins.




Figure 5. Phosphorylation of bovine rhodopsin by GRK7 and GRK1. Extracts from nontransfected (NT) HEK-293 cells or cells expressing bovine GRK1 or ground squirrel GRK7 were used to phosphorylate bovine ROS membranes as described in the "Methods". ROS membranes containing bovine rhodopsin were incubated with [γ^{32} P]ATP in the presence (+) or absence (-) of light, then electrophoresed on a 10% SDS-polyacrylamide gel. Phosphorylation was visualized by phosphorimage analysis. The numbers at left represent protein molecular size markers (Biorad, Hercules, CA).

An interesting feature of this kinase is the presence of a C-terminal CAAX motif coding for the addition of a geranylgeranyl group to Cys-545. In contrast, GRK1 is farnesylated in every species examined, except chicken, where it is geranylgeranylated [21]. The functional consequence of this difference in isoprenylation may be predicted from studies of Inglese et al. [49], who reported that mutation of the C-terminal CAAX box of GRK1 from CVLS to CVLL, to promote geranylgeranylation instead of farnesylation, resulted in constitutive association with the plasma membrane. Therefore GRK7 may be more tightly associated with the membrane than GRK1. It is intriguing to speculate that this difference in posttranslational modification could contribute to the differences in rates of signal termination between rods and cones through differences in rates of phosphorylation. Further analysis of the biochemical properties of this kinase will be necessary to verify such a hypothesis.

Another feature shared by GRK7 with several members of the GRK family is its ability to be autophosphorylated. Thr-21 in GRK7 from the 13-lined ground squirrel is conserved with Ser-21 in bovine GRK1, but this site is absent in *O. latipes* GRK7, suggesting that it may not be critical for function. Both the 13-lined ground squirrel and *O. latipes* GRK7 possess a serine at a position corresponding to Ser-488 in bovine GRK1, but Thr-489 in GRK1, which is also autophosphorylated, is a glutamic acid in both GRK7 proteins. It has been suggested that phosphorylation of these residues in GRK1 enhances its release from phosphorylated rhodopsin, thereby serving to limit the stoichiometry of phosphorylation [50,51]. These residues may also play a role in ATP binding [46].

The existence of GRK7 in the retinas of the 13-lined ground squirrel, the eastern chipmunk and the pig, and the




Figure 6. Autophosphorylation of GRK7. Extracts from nontransfected HEK-293 cells (NT) and cells transfected with histidine-tagged GRK7 were incubated with Ni-NTA resin and washed to remove unbound protein (Flowthrough). Both the flowthrough and the protein bound to the resin were incubated with [γ^{32} P]ATP as described in the "Methods". The numbers at left represent protein molecular size markers (BioRad).

ability of rhodopsin to serve as a substrate for this kinase, raise several interesting questions: (1) Does this novel kinase play a role in mammalian cone visual signaling pathways, similar to the role played by GRK1 in rods, by phosphorylating the cone opsins? (2) If GRK7 participates in cone signaling, do differences exist in the kinetics of cone opsin phosphorylation that may account for the faster signal termination in cones? In the medaka fish, GRK7 was expressed only in cones and GRK1 only in rods [20]. However in mammals, it has been suggested that GRK1 is also in cones [21]. If GRK7 is also found in both rods and cones, it raises the interesting possibility that these two closely-related members of the GRK family cooperate in visual signaling.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. David Fenstermacher for advice and assistance with DNA sequence analysis software and Dr. Debora Farber for helpful discussions. This work was supported by NIH grants GM47438 (S.O.), GM43582 (E.R.W.), EY11498 (F.W.), and EY10573 (T.W.K), grants from The Foundation Fighting Blindness (S.O. and E.R.W.; F.W.) and a grant from Research to Prevent Blindness (F.W.).

REFERENCES

- Zhao X, Palczewski K, Ohguro H. Mechanism of rhodopsin phosphorylation. *Biophys Chem* 1995; 56:183-8.
- Nathans J, Thomas D, Hogness DS. Molecular genetics of human color vision: the genes encoding blue, green, and red pigments. *Science* 1986; 232:193-202.
- Lochrie MA, Hurley JB, Simon MI. Sequence of the alpha subunit of photoreceptor G protein: homologies between transducin, ras, and elongation factors. *Science* 1985; 228:96-9.
- Lerea CL, Somers DE, Hurley JB, Klock IB, Bunt-Milam AH. Identification of specific transducin alpha subunits in retinal rod and cone photoreceptors. *Science* 1986; 234:77-80.
- Levine MA, Smallwood PM, Moen PT Jr, Helman LJ, Ahn TG. Molecular cloning of beta 3 subunit, a third form of the G protein beta-subunit polypeptide. *Proc Natl Acad Sci U S A* 1990; 87:2329-33.
- Ong OC, Yamane HK, Phan KB, Fong HK, Bok D, Lee RH, Fung BK. Molecular cloning and characterization of the G protein gamma subunit of cone photoreceptors. *J Biol Chem* 1995; 270:8495-500.
- Li TS, Volpp K, Applebury ML. Bovine cone photoreceptor cGMP phosphodiesterase structure deduced from a cDNA clone. *Proc Natl Acad Sci U S A* 1990; 87:293-7.
- Viczian AS, Piriev NI, Farber DB. Isolation and characterization of a cDNA encoding the alpha' subunit of human cone cGMP-phosphodiesterase. *Gene* 1995; 166:205-11.
- Shimizu-Matsumoto A, Itoh K, Inazawa J, Nishida K, Matsumoto Y, Kinoshita S, Matsubara K, Okubo K. Isolation and chromosomal localization of the human cone cGMP phosphodiesterase gamma cDNA (PDE6H). *Genomics* 1996; 32:121-4.
- Craft CM, Whitmore DH, Wiechmann AF. Cone arrestin identified by targeting expression of a functional family. *J Biol Chem* 1994; 269:4613-9.
- Craft CM, Whitmore DH. The arrestin superfamily: cone arrestins are a fourth family. *FEBS Lett* 1995; 362:247-55.
- Bonigk W, Muller F, Middendorff R, Weyand I, Kaupp UB. Two alternatively spliced forms of the cGMP-gated channel alpha subunit from cone photoreceptor are expressed in the chick pineal organ. *J Neurosci* 1996; 16:7458-68.
- Yau KW. Phototransduction mechanism in retinal rods and cones. The Friedenwald Lecture. *Invest Ophthalmol Vis Sci* 1994; 35:9-32.
- Imai H, Imamoto Y, Yoshizawa T, Shichida Y. Difference in molecular properties between chicken green and rhodopsin as related to the functional difference between cone and rod photoreceptor cells. *Biochemistry* 1995; 34:10525-31.
- Imai H, Kojima D, Oura T, Tachibanaki S, Terakita A, Shichida Y. Single amino acid residue as a functional determinant of rod and cone visual pigments. *Proc Natl Acad Sci U S A* 1997; 94:2322-6.
- Cowan CW, Fariss RN, Sokal I, Palczewski K, Wensel TG. High expression levels in cones of RGS9, the predominant GTPase accelerating protein of rods. *Proc Natl Acad Sci U S A* 1998; 95:5351-56.
- He W, Cowan CW, Wensel TG. RGS9, a GTPase accelerator for phototransduction. *Neuron* 1998; 20:95-102.
- Premont RT, Inglese J, Lefkowitz RJ. Protein kinases that phosphorylate activated G protein-coupled receptors. *FASEB J* 1995; 9:175-82.
- Palczewski K. GTP-binding-protein-coupled receptor kinases—two mechanistic models. *Eur J Biochem* 1997; 248:261-9.
- Hisatomi O, Matsuda S, Satoh T, Kotaka S, Imanishi Y, Tokunaga F. A novel subtype of G-protein-coupled receptor kinase, GRK7, in teleost cone photoreceptors. *FEBS Lett* 1998; 424:159-64.
- Zhao X, Huang J, Khani SC, Palczewski K. Molecular forms of human rhodopsin kinase (GRK1). *J Biol Chem* 1998; 273:5124-31.
- Yamamoto S, Sippel KC, Berson EL, Dryja TP. Defects in the rhodopsin kinase gene in the Oguchi form of stationary night blindness. *Nat Genet* 1997; 15:175-8.
- Khani SC, Nielsen L, Vogt TM. Biochemical evidence for pathogenicity of rhodopsin kinase mutations correlated with the Oguchi form of congenital stationary night blindness. *Proc Natl Acad Sci U S A* 1998; 95:2824-7.
- Cideciyan AV, Zhao X, Nielsen L, Khani SC, Jacobson SG, Palczewski K. Null mutation in the rhodopsin kinase gene slows recovery kinetics of rod and cone phototransduction in man. *Proc Natl Acad Sci U S A* 1998; 95:328-33.
- Cassill JA, Whitney M, Joazeiro CA, Becker A, Zuker CS. Isolation of Drosophila genes encoding G protein-coupled receptor kinases. *Proc Natl Acad Sci U S A* 1991; 88:11067-70.
- Haribabu B, Snyderman R. Identification of additional members of human G-protein-coupled receptor kinase multigene family. *Proc Natl Acad Sci U S A* 1993; 90:9398-402.
- Kunapuli P, Benovic JL. Cloning and expression of GRK5: a member of the G protein-coupled receptor kinase family. *Proc Natl Acad Sci U S A* 1993; 90:5588-92.
- Premont RT, Koch WJ, Inglese J, Lefkowitz RJ. Identification, purification, and characterization of GRK5, a member of the family of G protein-coupled receptor kinases. *J Biol Chem* 1994; 269:6832-41.
- Neill JD, Musgrove LC, Sellers JC, Duck LW. Identification by polymerase chain reaction of the G protein-coupled receptor kinases expressed in the mouse gonadotropic alpha T3-1 cell line. *Endocrinology* 1996; 137:3942-7.
- Frohman MA, Dush MK, Martin GR. Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proc Natl Acad Sci U S A* 1988; 85:8998-9002.
- Levison SW, Ducceschi MH, Young GM, Wood TL. Acute exposure to CNTF in vivo induces multiple components of reactive gliosis. *Exp Neurol* 1996; 141:256-68.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227:680-5.

33. Osawa S, Weiss ER. The effect of carboxyl-terminal mutagenesis of G_i alpha on rhodopsin and guanine nucleotide binding. *J Biol Chem* 1995; 270:31052-8.
34. Benovic JL, Gomez J. Molecular cloning and expression of GRK6. A new member of the G protein-coupled receptor kinase family. *J Biol Chem* 1993; 268:19521-7.
35. Kelleher DJ, Johnson GL. Characterization of rhodopsin kinase purified from bovine rod outer segments. *J Biol Chem* 1990; 265:2632-9.
36. Shi W, Osawa S, Dickerson CD, Weiss ER. Rhodopsin mutants discriminate sites important for the activation of rhodopsin kinase and G_i. *J Biol Chem* 1995; 270:2112-9.
37. Hara S, Miyoshi M. Gap junctions between visual cells in the squirrel retina. *Biomedical Research* 1984; 5:101-10.
38. von Schantz M, Szel A, van Veen T, Farber DB. Expression of phototransduction cascade genes in the ground squirrel retina. [published erratum appears in *Invest Ophthalmol Vis Sci* 1995; 36:3] *Invest Ophthalmol Vis Sci* 1994; 35:2558-66.
39. von Schantz M, Szel A, van Veen T, Farber DB. Expression of soluble phototransduction-associated proteins in ground squirrel retina. *Invest Ophthalmol Vis Sci* 1994; 35:3922-30.
40. Tansley K. Comparative anatomy of the mammalian retina with respect to the electroretinographic response to light. In: Smelser GK, editor. *The structure of the eye*. New York: Academic Press; 1961. p. 193-206.
41. Prince JH, Diesem CD, Eglitis I, Ruskell GL. Anatomy and histology of the eye and orbit in domestic animals. Springfield (IL): C. C. Thomas; 1960. pp. 210-33.
42. Gerke CG, Hao Y, Wong F. Topography of rods and cones in the retina of the domestic pig. *Hong Kong Medical Journal* 1995; 1:302-8.
43. Premont RT, Macrae AD, Stoffel RH, Chung N, Pitcher JA, Ambrose C, Inglese J, MacDonald ME, Lefkowitz RJ. Characterization of the G protein-coupled receptor kinase GRK4. Identification of four splice variants. *J Biol Chem* 1996; 271:6403-10.
44. Sallèse M, Mariggio S, Collodel G, Moretti E, Piomboni P, Baccetti B, De Blasi A. G protein-coupled receptor kinase GRK4. Molecular analysis of the four isoforms and ultrastructural localization in spermatozoa and germinal cells. *J Biol Chem* 1997; 272:10188-95.
45. Palczewski K, Buczylo J, Van Hooser P, Carr SA, Huddleston MJ, Crabb JW. Identification of the autophosphorylation sites in rhodopsin kinase. *J Biol Chem* 1992; 267:18991-8.
46. Palczewski K, Ohguro H, Premont RT, Inglese J. Rhodopsin kinase autophosphorylation. Characterization of site-specific mutations. *J Biol Chem* 1995; 270:15294-8.
47. Kunapuli P, Gurevich VV, Benovic JL. Phospholipid-stimulated autophosphorylation activates the G protein-coupled receptor kinase GRK5. *J Biol Chem* 1994; 269:10209-12.
48. Loudon RP, Benovic JL. Expression, purification, and characterization of the G protein-coupled receptor kinase GRK6. *J Biol Chem* 1994; 269:22691-97.
49. Inglese J, Koch WJ, Caron MG, Lefkowitz RJ. Isoprenylation in regulation of signal transduction by G-protein-coupled receptor kinases. *Nature* 1992; 359:147-50.
50. Buczylo J, Gutmann C, Palczewski K. Regulation of rhodopsin kinase by autophosphorylation. *Proc Natl Acad Sci U S A* 1991; 88:2568-72.
51. Pulvermüller A, Palczewski K, Hofmann KP. Interaction between photoactivated rhodopsin and its kinase: stability and kinetics of complex formation. *Biochemistry* 1993; 32:14082-8.

The typographical corrections below were made to the article on the date noted. These changes have been incorporated in the article and the details are documented here.

9 December 1998: In paragraph 6 of Results, "Figure 1" was changed to "Figure 2".

16 December 1998: In paragraphs 7, 12 and 13 of Methods and the Figure 5 caption, "chromatographed" was changed to "electrophoresed."